Issued May 27, 1910.

Return to Forage Crop Investigations files.

U. S. DEPARTMENT OF AGRICULTURE.

FARMERS' BULLETIN 400.

# A MORE PROFITABLE CORN-PLANTING METHOD.

BY

C. P. HARTLEY,

*Physiologist in Charge of Corn Investigations, Bureau of Plant Industry.*

WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1910.

# LETTER OF TRANSMITTAL.

U. S. DEPARTMENT OF AGRICULTURE,
BUREAU OF PLANT INDUSTRY,
OFFICE OF THE CHIEF,
*Washington, D. C., April 1, 1910.*

SIR: I have the honor to transmit herewith, and to recommend for publication as a Farmers' Bulletin, a manuscript entitled "A More Profitable Corn-Planting Method," by C. P. Hartley, Physiologist in Charge of Corn Investigations.

Studies of the habits of growth both below and above ground of this our most valuable and widely grown plant have for several years indicated that fields of checked corn, comprising about half of the total corn acreage, yield less than they would if the kernels were not crowded so closely together in the hills.

As this assumption has now been substantiated, the farmers should be informed on the subject and advised to substitute, as soon as possible, some method of checking that will give every stalk in their cornfields a fair chance to thrive and produce abundantly.

Respectfully,

G. H. POWELL,
*Acting Chief of Bureau.*

Hon. JAMES WILSON,
*Secretary of Agriculture.*

# CONTENTS.


| | Page. |
|---|---|
| Introduction | 5 |
| Methods by which most corn is planted | 5 |
| The advantage of drilling | 5 |
| The advantage of checking | 5 |
| Double advantages of kernel-spaced checking | 6 |
| Why kernel-spaced checking should give increased yields | 8 |
| Kernel-spaced checking tested in comparison with the usual method | 8 |
| Kernel-spaced checking yields better in every case | 11 |
| The increased yield is net gain | 11 |
| Kernel-spaced checking permits cultivation without difficulty | 12 |
| Other advantages of kernel spacing | 12 |
| Present application of the advantages | 13 |
| Profit depends mainly on large yields to the acre | 13 |
| Summary | 14 |


# ILLUSTRATIONS.


| | | Page. |
|---|---|---|
| FIG. 1. | Hill of corn planted by a modern check-rower | 6 |
| 2. | Base of a corn plant at silking time | 7 |
| 3. | The device used in dropping corn by hand in the kernel-spaced rows of the tests | 8 |
| 4. | A hill of the kernel-spaced rows of the tests | 9 |
| 5. | A hill of the rows tested in alternation with the kernel-spaced rows | 9 |




400

B. P. I.—571.

# A MORE PROFITABLE CORN-PLANTING METHOD.

## INTRODUCTION.

Much study has been given to corn-planting methods and much progress made. This progress has usually been along the line of more rapid and easier planting and cultivating. While retaining these advantages, the method here described causes higher yields by supplying each plant with better conditions for growth and production.

## METHODS BY WHICH MOST CORN IS PLANTED.

Single-row corn planters drawn by one horse—or two-row planters drawn by two horses—are used in planting the greater part of the corn grown in the United States. All these planters deposit the kernels of hill-planted corn together in a small space, or bunch, resulting in the stunting of many plants because of insufficient space to form a strong root system or vigorous stalk and leaves. Many prevent this stunting by drilling their corn. Drilled corn is planted in rows, one kernel in a place. Checked corn is planted in hills that permit cultivation both ways.

Which is the most profitable, drilling or checking, is a much debated question. On hilly land, where the soil washes readily, checking is not practicable. Where both methods are practicable one method has about as many advantages and disadvantages as the other, and corn producers are practicing the two methods to about the same extent.

## THE ADVANTAGE OF DRILLING.

Standing singly, as in drilled corn, each plant has a fairer chance both below ground and above ground to develop normally and produce well.

It requires a little less care to drill than to check, though in fields free from trees the difference is very slight after the operator has become accustomed to the use of the checking wire which is stretched across the field.

## THE ADVANTAGE OF CHECKING.

Checking enables the corn to be cross-cultivated and kept free from weeds and the entire soil surface kept in good condition without the expensive labor of hoeing.



400

## DOUBLE ADVANTAGES OF KERNEL-SPACED CHECKING.

"Kernel-spaced checking" is the term used to designate the more profitable method described in this bulletin. It largely overcomes the disadvantages and combines the advantages of the two methods now used in planting most of the corn grown in the United States.



FIG. 1.—Hill of corn planted by a modern check-rower. The central stalk had no chance to develop, either below or above ground, and produced but a nubbin.

The double advantages of kernel-spaced checking became apparent several years ago in connection with tests of thick planting on very fertile river-bottom soil. On this soil one stalk to every 12 inches of row, when hoed, was found to give higher yields than checked corn with either three or four stalks in every hill.

Drilling, however, was not practicable because of the abundant growth of weeds. Planting four stalks in each hill with a check-rower was found to result in an unusually large percentage of nubbins and stunted and barren stalks. Drilled corn with the same number of stalks to the acre gave larger yields with fewer nubbins and stunted and barren stalks, but could not be well cultivated without expensive hoeing.

It was thought that kernel-spaced checking would be found profitable on soil sufficiently well supplied with moisture and fertility to permit four or five stalks to the hill. This has been found to be the case. Furthermore, it has been found profitable on dry soils of medium fertility.

In all fields of corn planted with our most modern check-rowers can be found many hills containing a stalk that has not produced

FIG. 2.—Base of a corn plant at silking time. For a foothold each plant needs a cubic foot of space.

well. In many such cases the stunting is due to the quicker starting or better location of the other plants, which enabled them to monopolize the space, moisture, fertility, and sunlight of the limited area in which the kernels were planted. (See fig. 1.)

With roots and stalks limited to so small a space, plants from any cause once obtaining a monopoly easily retain it. This plausible idea is substantiated by the tests described later, which produced

more stunted and barren stalks and more nubbins in rows of ordinary checking than in adjacent rows of kernel-spaced checking.

Such stunting, because of the nature of the corn plant, materially reduces the grain yield. Plants that are crowded to such an extent that they become slender and scarcely more than half their normal height are either barren or produce only nubbins.

## WHY KERNEL-SPACED CHECKING SHOULD GIVE INCREASED YIELDS.

The disadvantage under which the two or more plants of a hill are compelled to grow by the kernels having been bunched together is so plain that it is remarkable we have not long ago devised a method of checking corn that gives each plant room to form a normal root system.



FIG. 3.—The device used in dropping corn by hand in the kernel-spaced rows of the tests.

For a distance of 6 inches in every direction from the base of a normal corn plant the soil is crowded entirely full of roots. When several kernels are bunched together the roots of all the stalks of the hill are so matted together that the pulling of one stalk pulls out the entire hill, and the pressure of wind upon all the stalks is more likely to blow over the entire hill than when the stalks stand several inches apart in the hill, thus forming a broader base of support. Figure 2 shows the base of one corn plant at silking time, the critical period, when all the roots should be well located for supplying moisture to the plant. The three stalks of a hill grow at a great disadvantage with no more foothold in the soil than one plant requires.

## KERNEL-SPACED CHECKING TESTED IN COMPARISON WITH THE USUAL METHOD.

The following-described tests were especially planned to decide whether the kernel spacing of checked corn would increase the yield above that obtained by the generally practiced method of checking. Eight alternating rows of kernel-spaced and of ordinary checking were planted in three different localities on different types of soil. In order to test the practicability of cultivating the kernel-spaced checking, the test rows were placed in the center of fields of checked corn and were without difficulty cultivated both ways when the fields were cultivated.

The device used for space planting is shown in figure 3. Two kernels were planted at each corner of a 5-inch square. Each space-



FIG. 4.—A hill of the kernel-spaced rows of the tests. The stalks stand at the corners of a 5-inch square. (Photographed before the formation of brace roots at the lower nodes.)

planted row was followed by a row treated in all respects the same except that the kernels were not so well separated, being deposited as a check-rower would deposit them and with six kernels to the hill.

FIG. 5.—A hill of the rows tested in alternation with the kernel-spaced rows. These hills were bunched to no greater extent than is done by the modern check-rower. (See fig. 1.)

When the corn was a foot tall, all the rows were thinned to a uniform stand. Four stalks to the hill were retained wherever possible. When about ready to tassel several more stalks were removed from some of the rows to bring adjacent rows to exactly the same number of stalks.

Figures 4 and 5 show the differences in appearance due to the different methods of

planting. Figure 4 shows a typical hill from the space-planted rows, while figure 5 is typical of the usual method of checking. The difference in the uniformity of the stalks is very marked.

TABLE I.—*Comparative yields of corn when kernel-spaced checking and the usual method of checking were employed.*

| | Tests at Round Hill, Va. | Tests at Piketon, Ohio. | Tests at McLean, Va. |
|---|---|---|---|
| Character of the summer | Very dry | Very favorable | Very dry. |
| Nature of the soil | Stony clay loam. | River-bottom deposit. | Heavy clay loam. |
| Distances between corn hills | 3¾ by 3⅓ feet. | 3⅓ by 3¼ feet | 3¾ by 3½ feet. |
| Height of the varieties tested | Tall | Very tall | Medium. |
| Average number of stalks to the hill. | 3 | 3¾ | 3⅘. |
| Number of stalks to the acre | 10,500 | 15,100 | 12,700. |
| Stalk circumference increased by spacing. | 14½ per cent. | 7 per cent | 13 per cent. |
| Number of poor ears produced: | | | |
| Kernel-spaced method | 498 | 447 | 652. |
| Usual method | 584 | 502 | 712. |
| Yield per acre: | | | |
| Kernel-spaced method | 55 bushels | 118½ bushels | 64 bushels. |
| Usual method | 53 bushels | 114 bushels | 61 bushels. |
| Gain to the acre by spacing the kernels. | 2 bushels | 4½ bushels | 3 bushels. |
| Number of adjacent-row tests | 7 | 9 | 9. |
| Number of times in which adjacent-row tests produced equally. | 0 | 1 | 1. |
| Number of tests in which kernel spacing increased the yields. | 4 | 7 | 8. |
| Increase obtained by spacing the kernels. | 3¾ per cent | 4 per cent | 5 per cent. |

## KERNEL-SPACED CHECKING YIELDS BETTER IN EVERY CASE.

Under the different conditions of climate and soil and with different varieties of corn, each test has shown that kernel spacing of checked corn increases the yield. The kernels were placed only 5 inches apart in the hills. A distance of 8 or 10 inches may increase the yield still more, and such hills can be readily cultivated both ways provided the checking is accurately done. Check-rowers can be manufactured with two shoes for each corn row and a separate dropping tube for each kernel of a hill. They can be made adjustable, so that the number of kernels dropped and the spacing distances between the kernels of a hill can be suited to the fertility of the soil and the variety of corn planted.

Proper distances for spacing the kernels within the hill will have to be determined for large and small growing varieties. The three varieties tested ranged from medium to very tall, each being adapted to the locality in which it was tested. All produced better by spacing the kernels of each hill 5 inches apart than by the customary method of dropping them close together in the hill.

In addition to the 25 tests of rows growing side by side in which one row was kernel-spaced and the adjacent row checked by the usual method, there were at each locality two adjacent plats, one kernel-spaced checked and the other checked by the usual method. At two of the localities the yield of the kernel-spaced plats exceeded that of those planted by the usual method to about the same extent as in the alternate-row tests. At McLean, Va., the plats produced equally well, but the alternate-row plantings gave an increase of 11 per cent in yield due to kernel spacing, bringing the average increase for the two methods of testing to practically 5 per cent, as given in the table.

## THE INCREASED YIELD IS NET GAIN.

Two bushels more to the acre, or a 4 per cent increase, is not small enough to be ignored, especially when obtainable without cost except the initial cost of a kernel-spacing check-rower.

Increased yields resulting from applications of expensive fertilizers sometimes fail to increase the profit on the crop. The increased yield due to spacing the kernels is clear gain, since a properly constructed check-rower can drop the seed in separate places as rapidly and as cheaply as the present check-rower bunches them.

In planting 50 acres of corn such a check-rower would pay for itself the first year. In planting 100 acres, estimating the production at the average of 26 bushels, it would return a net profit of 104 bushels.

A 4 per cent increase in the value of the corn crop of Illinois for one year would amount to $6,000,000. About one-half of the corn acreage of the United States is checked or planted with planters that drop all the kernels of a hill in a bunch. The substitution of kernel-spacing planters would add 50,000,000 bushels of corn to our annual production.

## KERNEL-SPACED CHECKING PERMITS CULTIVATION WITHOUT DIFFICULTY.

The 5-inch spacing of the kernels in each hill did not make it difficult to properly cultivate in both directions with 2-horse double cultivators, although in one instance the rows were but 3⅓ feet apart in one direction and 3¼ feet in the other. Three-horse 2-row cultivators could have been used to as good advantage as in the usual method of checking. If the kernels in each hill are spaced 6 or 8 inches apart, the gangs of shovels on the cultivator will need to be set farther apart in order to properly straddle the hills. A spacing wider than 8 inches may allow weeds to grow in the corn hills. In this respect the 5-inch kernel spacing gave no greater growth of weeds and received no more work than the ordinary checking.

## OTHER ADVANTAGES OF KERNEL SPACING.

Increased yield without extra cost is abundant reason to cause a change of methods. A few other advantages of kernel spacing may be mentioned. It is easier to thin kernel-spaced hills, and the thinning can be done without injury to the remaining stalks.

When the plants stand separately in the hills, an entire hill is less liable to be pulled out by the harrow, the weeder, or the cultivator. The entire hill is not so likely to be injured by accident, disease, cutworms, or corn-root insects. The better distribution not only affords each stalk better sunlight, but also shades the soil better, thereby lessening evaporation from the heated soil during the summer months.

At McLean, Va., the rows checked by the usual method blew down at tasseling time to a much greater extent than the adjacent kernel-spaced rows. At Piketon, Ohio, the kernel-spaced rows were noticed at harvest time to have fewer down hills than the adjacent bunch-planted rows. The kernel-spaced stalks appeared stronger and measurements proved them to be at each locality of greater circumference than the bunch-planted stalks. (See Table I.)

## PRESENT APPLICATION OF THE ADVANTAGES.

Until kernel-spacing check-rowers are placed upon the market farmers can utilize this method extensively only by remodeling their check-rowers or by double driving. By returning across the field with the check-rower, without resetting the wire, and driving 6 inches from the previous track, each hill can be made up of two bunches of kernels. This will improve somewhat the distribution of the plants in the hills, but will not be as good as a check-rower with four or more dropping tubes for each corn row.

By double driving, as just mentioned, with planters set to drop one kernel at a time, each stalk will be given room to form a good root system. Drilled corn planted by this method won the Iowa sweepstakes prize in 1909 by producing 153 bushels of shelled corn on 1 acre.[a]

## PROFIT DEPENDS MAINLY ON LARGE YIELDS TO THE ACRE.

Increasing population and increasing value of land are sure to stop the long-continued practice of planting corn thinner as the land becomes poorer, making up the supply by planting more acres. Our future increase in corn production must be accomplished mainly by filling the soil with water-absorbing and water-holding humus, accompanied by thicker planting. This practice has produced 239 bushels of shelled corn on an acre in South Carolina,[b] and 226 bushels on an acre in North Carolina.[c]

Soils are not impoverished by producing large crops. On the contrary, the larger the crops and the more frequently they are produced the better the opportunity to enrich the soil by plowing under a sufficient portion of the vegetable matter produced. The soil should be kept busy enriching itself through the decay of vegetation, as forest lands enrich themselves. To most soils one or more mineral elements must eventually be applied to replace those which are removed from the farm in the form of grain or animal products.

---

[a] See Farmers' Tribune, February 3, 1910. John Sundberg, of Monona County, Iowa, won this prize in accordance with the rules of the Iowa Corn Growers' Association. The planter was driven the second time several inches from its previous track, giving a double row of plants. Adjusted to drop one kernel every 16 inches the planter driven in this manner planted 20,633 kernels to the acre. The acre had been a clover pasture for five years and was fall-plowed. No manure or commercial fertilizer was used.

[b] See C. S. Plumb's book on Indian Corn Culture, 1895, p. 220. The rows were alternately 3 and 6 feet apart, and 5 or 6 kernels were dropped to each foot of the row, 1,000 bushels of stable manure, 600 bushels of cotton seed, and $50 worth of commercial fertilizer being applied to the acre.

[c] See Wake County Corn Experiments of 1909, by F. A. Olds, secretary, Chamber of Commerce, Raleigh, N. C. The first prize was won by J. F. Batts, of Garner, N. C. The rows were 41 inches apart, and 3 kernels were planted in hills 8 inches apart in the row, 19,000 stalks being grown on 1 acre. Forty-five 2-horse loads of manure and $59 worth of commercial fertilizer were applied to the acre.

## SUMMARY.

In planting corn in hills practically all corn planters drop all the kernels of the hill in a bunch.

A study of the best conditions for growth and production indicates that bunch planting results in unnecessary crowding both above and below the ground, weakening the stalks and reducing the yield.

Kernel-spaced checking combines the advantages and overcomes the disadvantages of drilling and checking, the two methods by which nearly all corn is now planted.

Tests with three varieties of corn under different conditions of soil and climate in every case gave increased yields due wholly to a separation of the kernels planted in each hill.

# INDEX.


| | Farmers' Bull. No. | Page. |
|---|---|---|
| Acetanilid, character and use | 377 | 3 |
| harmful effects, investigation, results, summary | 377 | 6–7 |
| introduction as medicine | 377 | 3 |
| poisoning, number of cases, symptoms, fatalities, etc | 377 | 9–13 |
| use as habit-forming agent | 393 | 3 |
| dose, harmfulness, etc., reports of physicians | 377 | 6–9 |
| objects | 377 | 4–5 |
| Acetphenetidin, use as habit-forming agent | 393 | 3 |
| *Achroria grisella*. *See* Wax moth, lesser. | | |
| Acid, citric, experiments in jelly making | 388 | 31–32 |
| development in butter and cheese making, amount | 381 | 32 |
| tartaric, experiments in jelly making | 388 | 31–32 |
| Acidimeter, use in making butter and cheese, description | 381 | 30–31 |
| Acidity, control in butter and cheese making, study of methods and apparatus | 381 | 30–32 |
| measuring, method in manufacturing Camembert cheese | 384 | 26 |
| Acids, experiments in jelly making | 388 | 31–32 |
| Adulterants, seed, testing method | 382 | 21–23 |
| Adulteration, forage plant seeds | 382 | 1–23 |
| honey, laws affecting, note | 397 | 42 |
| Agricultural clubs, boys' and girls' | 385 | 1–23 |
| junior, available publications, list | 385 | 19–22 |
| organization and work, suggestions | 385 | 13–19 |
| statistics | 385 | 22–23 |
| college, Cornell University, boys' and girls' clubs, responsibility | 385 | 7 |
| Department, cooperation in boys' and girls' argicultural club work | 385 | 10–11 |
| Air-sac mite, disease affecting adult pheasants | 390 | 39 |
| Alfalfa, loco poisoning antidote | 380 | 13 |
| meal, grinding, study of expenses and benefits, and results | 384 | 13–14 |
| protein content, comparison with timothy hay | 384 | 13 |
| wheat bran, experiments in Pennsylvania | 384 | 13–14 |
| value as cow feed, comparison with wheat bran, experiments in Vermont and Massachusetts | 384 | 14 |
| stock feed, study and experiments | 384 | 12–14 |
| rain requirements in various States | 388 | 11 |
| relation of yellow trefoil | 382 | 5 |
| seed, adulteration with sweet clover | 382 | 9 |
| yellow trefoil | 382 | 8 |
| use as legume in potato culture | 386 | 5 |
| Alkali, presence in swill, responsibility for hog disease | 379 | 15–16 |
| Animals, farm. *See* Farm animals. | | |
| Anthracnose, fungous disease of bean | 388 | 21–23 |
| Antidotes, loco weed poisoning | 380 | 13–14 |
| Antipyrin, character and use | 377 | 3 |
| harmful effects, investigation, results, summary | 377 | 6–7 |
| introduction as medicine | 377 | 3 |
| poisoning, number of cases, symptoms, fatalities, etc | 377 | 13–14 |
| use as habit-forming agent | 393 | 3 |
| dose, harmfulness, etc., reports of physicians | 377 | 6–9 |
| objects | 377 | 4–5 |
| Apiary, beginning, purchasing bees, selection, etc | 397 | 11–13 |
| location, equipment, etc | 397 | 6–13 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| *Aragallus lamberti*, description | 380 | 7–8 |
| *Argusianus*, pheasant variety | 390 | 10 |
| Arid lands, west, use of windmills in irrigation | 394 | 1–44 |
| Arid-land farming. *See* Dry farming. | | |
| Arizona, early onion raising, experiments and results | 384 | 6, 7, 8 |
| Arsenic, loco poisoning antidote | 380 | 13 |
| use in cattle dip, precautions | 378 | 24 |
| Asthma remedies, harmful nature, notes | 393 | 8–10 |
| *Astragalus arizonicus*, studies | 380 | 9 |
| *bigelovii*, studies | 380 | 9 |
| *diphysus*, studies | 380 | 9 |
| *mollissimus*, description | 380 | 6–7 |
| *thurberi*, studies | 380 | 9 |
| ATWATER, HELEN W., bulletin on "Bread and bread making" | 389 | 1–47 |
| Auger, stump, description, etc | 381 | 6–7 |
| Austria, artificial feeds for calves, experiments and results | 381 | 21–22 |
| Bache, Richard, early importation of pheasants and partridges | 390 | 13 |
| *Bacillus mesentericus vulgatus*, effect on bread | 389 | 34 |
| Bacteria, lactic-acid, use in manufacture of skim-milk buttermilk | 384 | 18–19 |
| Baking bread, methods | 389 | 28–30 |
| Banat bees, note | 397 | 13 |
| Barium, presence in loco weed | 380 | 8–9, 13 |
| Barley flour, use in bread making, note | 389 | 14 |
| tillage methods, experiments in Montana | 388 | 8 |
| Barm, use in bread making | 389 | 24–25 |
| Bean, moth, value for hay production | 388 | 11 |
| soup, use in feeding calves without milk | 381 | 15 |
| Beans, anthracnose, origin, control methods, etc | 388 | 21–23 |
| BEAVERS, J. C., bulletin on "Farm practice in the use of commercial fertilizers in the South Atlantic States" | 398 | 1–24 |
| Bee colony, description, habits, characteristics, etc | 397 | 14–17 |
| queenless, management | 397 | 26 |
| diseases, laws affecting, note | 397 | 42 |
| hives, types, selection, etc | 397 | 8–10 |
| keepers' associations, notes | 397 | 41 |
| keeping, directions, apparatus, laws, etc | 397 | 1–44 |
| publications | 397 | 43–44 |
| "Beef birds," recipe | 391 | 27 |
| different cuts, proportion, relation to prices, etc | 391 | 8–10 |
| Bees, behavior, necessity for study of, by bee keepers | 397 | 14 |
| studies | 397 | 14–17 |
| care, management, diseases, etc | 397 | 1–44 |
| different races, characteristics, etc | 397 | 12–13 |
| diseases | 397 | 38–39 |
| enemies | 397 | 40 |
| habits, necessity for knowledge of, by bee keepers | 397 | 14 |
| manipulation, directions | 397 | 17–26 |
| queen, buying, suggestions | 397 | 12–13 |
| robbing other colonies, prevention | 397 | 23 |
| swarming, definition and description | 397 | 15 |
| transferring, directions | 397 | 20–22 |
| wintering, preparation | 397 | 36–38 |
| Beeswax, production | 397 | 36 |
| Beet crop, failures under irrigation, remedies, etc | 392 | 40 |
| culture under irrigation | 392 | 28–38 |
| puller, description, illustration, etc | 392 | 34 |
| Beets, sugar, cost of growing and returns | 392 | 44–45 |
| industry, historical notes, etc | 392 | 7–9 |
| irrigation | 392 | 52 |
| results of experiments | 392 | 45–47 |
| Beet-sugar factories, number and capacities by States, 1908 | 392 | 8 |
| Belling, J., experiments in jelly making from guava juice | 388 | 29 |
| Biscuits, Maryland, making | 389 | 26–27 |
| Black bees, remarks | 397 | 12 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| Blocking sugar beets | 392 | 31–32 |
| Blood, dried, cost, comparison with cotton-seed meal | 388 | 25 |
| use as feed for farm stock | 388 | 24–25 |
| Bluegrass, Canada, relation to Kentucky bluegrass | 382 | 5 |
| Kentucky, relation of Canada bluegrass | 382 | 5 |
| Bonavist, drought resistance | 388 | 11 |
| Bones, uses in cooking | 391 | 20–21 |
| Boss, Andrew, statement on Kherson and Sixty-day oat tests in Minnesota | 395 | 18 |
| Braised beef, recipe | 391 | 29–30 |
| Bran, wheat, comparison with alfalfa meal as cow feed, experiments in Vermont and Massachusetts | 384 | 14 |
| protein content, comparison with alfalfa meal, experiments in Pennsylvania | 384 | 13–14 |
| Bread, aerated, making | 389 | 20–21 |
| and bread making | 389 | 1–47 |
| chemical composition | 389 | 36–37 |
| digestibility of different kinds | 389 | 42–45 |
| food value, early use, etc | 389 | 5–7 |
| imperfections, causes and remedies | 389 | 33–35 |
| making and bread | 389 | 1–47 |
| losses of material | 389 | 33 |
| place in diet | 389 | 45–46 |
| relation of quality to gluten of the flour | 389 | 31–32 |
| "salt-rising," making | 389 | 21 |
| stale, notes | 389 | 31 |
| unleavened, making | 389 | 27–28 |
| Breads, composition, compared with other foods | 389 | 37–40 |
| Brewster, C. E., T. S. Palmer, and Henry Oldys, bulletin on "Game laws for 1909" | 376 | 1–56 |
| Brome-grass, awnless, seed, adulteration, and adulterants | 382 | 14–15 |
| Brooders, colony, gasoline-heated, cost of operation, comparison with other brooders | 381 | 28–30 |
| description, use, cost, etc | 381 | 23–30 |
| Butter making, acid, development | 381 | 32 |
| acidity, measuring, method | 381 | 30–32 |
| farm, directions, conditions affecting, etc | 384 | 22–24 |
| discussion by F. Rasmussen | 384 | 22, 23 |
| markets, profits, etc | 384 | 22–24 |
| quality, price, etc., conditions affecting | 384 | 22–24 |
| Buttermilk, manufacture from skim milk, method | 384 | 18–19 |
| skim-milk, method of manufacture | 384 | 18–19 |
| Cacodylate, sodium, loco poisoning antidote | 380 | 14 |
| Cafe-Coca, harmful character, note | 393 | 8 |
| Caffein, use as habit-forming agent | 393 | 3 |
| Calf feeding, experiments and results | 381 | 14–23 |
| milk substitutes, experiments and results | 381 | 15 |
| California, experiments in pheasant propagation | 390 | 16 |
| windmill irrigation near Stockton | 394 | 41–44 |
| Calves, artificial feeds, experiments in various foreign countries, and results | 381 | 19–23 |
| feeding without milk, methods | 381 | 15 |
| raising in Illinois, cost | 381 | 17–18 |
| Massachusetts, cost | 381 | 17 |
| Canada, oat-seeding experiments, yields, etc | 388 | 13, 14, 15 |
| Cannabis indica, use in soothing sirup | 393 | 4–5 |
| Carniolan bees, note | 397 | 13 |
| Casserole cookery, recipes, methods, etc | 391 | 30–31 |
| Catarrh remedies, harmful nature, notes | 393 | 10–12 |
| Cattle, loco poisoning, symptoms | 380 | 11–12 |
| poisoning by loco weed | 380 | 9–11 |
| spraying for Texas-fever ticks, method | 378 | 24–27 |
| Caucasian bees, characteristics, value, etc | 397 | 13 |
| Celery Cola, harmful character, note | 393 | 8 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| Cellars, potato, method of building | 386 | 9 |
| Cement, use in manufacture of fence posts, method, etc | 384 | 29–32 |
| Center, O. D., statement on oat culture in Illinois | 395 | 16 |
| Cereals, analysis, table | 389 | 16 |
| use in bread making, etc | 389 | 8–16 |
| Charring, value in preservation of posts, methods | 387 | 7 |
| Checking, corn, advantages of various methods | 400 | 5–12 |
| kernel-spaced, double advantages of method | 400 | 6–12 |
| Check-rower, suggestions for manufacture and use in corn planting | 400 | 11–12, 13 |
| Cheese, Camembert, making room, equipment | 384 | 24–26 |
| manufacturing in United States | 384 | 24–29 |
| methods, possibilities, etc | 384 | 24–29 |
| making, acid development | 381 | 32 |
| acidity, measuring, method | 381 | 30–32 |
| Chess, use as adulterant in brome-grass | 382 | 14–15 |
| meadow fescue seed | 382 | 12 |
| Chickens, brooders, gasoline-heated, description, cost, etc | 381 | 23–30 |
| domesticated, descendants of jungle fowls | 390 | 10–11 |
| Chloral hydrate, use as habit-forming agent | 393 | 3 |
| in soothing sirups | 393 | 5 |
| Chloroform, use in soothing sirups | 393 | 4–5 |
| Cholera, disease affecting adult pheasants | 390 | 38 |
| hog. *See* Hog cholera. | | |
| *Chrysolophus amhersitæ*, pheasant variety | 390 | 8, 9 |
| *Cleonus quadrilineatus*, insect enemy of loco weeds | 380 | 15–16 |
| Clover, Bokhara. *See* Clover, sweet. | | |
| bur, use as adulterant in alfalfa seed | 382 | 8–9 |
| red, relation of yellow trefoil | 382 | 5 |
| seed, alsike, adulteration and adulterants | 382 | 10–11 |
| red, adulteration with screenings | 382 | 8 |
| yellow trefoil | 382 | 7 |
| sweet, use as adulterant in alfalfa seed | 382 | 9 |
| use as a legume in potato raising | 386 | 5 |
| Clubs, agricultural, boys' and girls' | 385 | 1–23 |
| State-wide movement, inception, growth, etc | 385 | 7–10 |
| various names, organization, work, etc | 385 | 5–10 |
| junior, available publications, list | 385 | 19–22 |
| organization, suggestions | 385 | 13–19 |
| statistics | 385 | 22–23 |
| Cornell Farm Boys' and Girls', organization, distribution, and total membership | 385 | 7 |
| Cocain, annual consumption | 393 | 3 |
| dangerous habit-forming drug | 393 | 6–7 |
| hydrochlorid. *See* Cocain. | | |
| use as habit-forming agent | 393 | 3 |
| danger to reformed addicts | 393 | 7 |
| Cocoa shells, decoction, use as substitute for skim milk, Canada | 381 | 19 |
| Codein, use as habit-forming agent | 393 | 3 |
| in soothing sirups | 393 | 5, 6 |
| Cold, remedies, harmful nature, notes | 393 | 12–13 |
| Colorado, experiments in pheasant propagation | 390 | 17 |
| mountain valleys, adaptability to potato culture | 386 | 5 |
| pruning experiments | 388 | 19–21 |
| tests of Sixty-day and Kherson oats | 395 | 24 |
| windmill irrigation | 394 | 39–41 |
| Comb honey, production for market, method | 397 | 32–36 |
| white, disease affecting pheasants | 390 | 39 |
| Concrete fence posts, manufacture, methods, cost, etc | 384 | 29–32 |
| mixer, home-made, description, use, etc | 384 | 30–31 |
| use in manufacture of concrete fence posts | 384 | 30–31 |
| Connecticut dairy cows, daily rations, cost, etc | 384 | 16–18 |
| herds, production, profit, etc., studies | 384 | 16–18 |
| Consumption cures, harmful nature, notes | 393 | 13–14 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| Corbett, L. C., bulletin on pruning, reference | 388 | 21 |
| Corn, checking, kernel-spaced, advantages, comparison with other methods, etc | 400 | 8–11 |
| drilling, advantages and disadvantages | 400 | 5, 6 |
| fertilizing, studies and suggestions | 398 | 19–21 |
| growing, responsibility of large yields for profit | 400 | 13 |
| sources of profit | 400 | 13 |
| planting, check-rower, disadvantages of use | 400 | 6, 7–8 |
| suggestions for manufacture and use | 400 | 11–12, 13 |
| kernel-spaced checking, double advantages of method | 400 | 6–12 |
| kernel-spacing, methods | 400 | 13 |
| methods, study and comparison | 400 | 5–12 |
| more profitable method | 400 | 1–14 |
| structure, use as food, etc | 389 | 15–16 |
| yield, comparison of checking methods | 400 | 8–11 |
| Corrugation irrigation, method, etc | 399 | 12–14 |
| Cotton, fertilizing, studies and suggestions | 398 | 17–19 |
| Cotton-seed meal, fermented, feeding to hogs, danger | 384 | 9–11 |
| Cough remedies, harmful nature, notes | 393 | 12–13 |
| Coverts, pheasant, necessity, description, etc | 390 | 17–19 |
| Cowpeas, fertilizing, studies | 398 | 23–24 |
| Cows, dairy, production, profit, etc | 384 | 16–18 |
| milch, value of mangels as feed | 384 | 14–15 |
| Cramps, disease affecting young pheasants | 390 | 37 |
| Cream, handling, care, etc., in farm butter making | 384 | 22–24 |
| importance as food, uses, care, etc | 384 | 19–22 |
| whipped, production, conditions affecting | 384 | 20–22 |
| uses, value as food, etc | 384 | 19–22 |
| whipping, methods, aids, etc | 384 | 19–22 |
| Creosote, application, methods, cooperative experiments, and results | 387 | 13–16 |
| impregnation, treating tanks, description | 387 | 10–11 |
| treatment, preparation of fence posts | 387 | 11–13 |
| various woods, experiments and results | 387 | 15–16 |
| use as shingle preservative, method, cost, etc | 387 | 18–19 |
| in preservation of farm timbers | 387 | 6, 7, 9–17 |
| Creosoting, fence posts, cost of treatment | 387 | 16 |
| value of treatment | 387 | 16–17 |
| plant, cooperative ownership, suggestions | 387 | 19 |
| Cresol solution, use as disinfectant in hog cholera prevention | 379 | 18 |
| Crop rotation, approved method | 388 | 9–10 |
| rotations, influence on use of commercial fertilizers | 398 | 6 |
| study of practices in South Atlantic States | 398 | 11 |
| Crops, injury by bees, supposed | 397 | 42 |
| under irrigation, production and value, estimates | 394 | 36 |
| various kinds, adaptability to various sections | 388 | 10–11 |
| winter cover, influence on use of commercial fertilizers | 398 | 7–8 |
| *Crossoptilon mantchuricum*, pheasant variety | 390 | 7, 8, 9–10 |
| Crowder, adjustable for making ditches for irrigation | 399 | 8 |
| Curbing, well, materials, etc | 394 | 10 |
| Curry, use in cooking meats | 391 | 38–39 |
| Cyprian bees, characteristics, note | 397 | 13 |
| Dairy herds, daily rations, cost, etc., studies in Connecticut | 384 | 16–18 |
| products, profits, etc., studies | 384 | 16–18 |
| records, importance of keeping | 384 | 16–18 |
| Dammer, use in irrigation ditches, description | 399 | 8–9 |
| Dams, irrigation ditches, description | 392 | 24–25 |
| Day, J. W., inventor of auger for use in removal of stumps | 381 | 6 |
| DEARBORN, NED, bulletin on "How to destroy English sparrows" | 383 | 1–11 |
| Delaware, experiments in pheasant propagation | 390 | 17 |
| Denny, O. N., introduction of pheasants into Oregon | 390 | 14 |
| Diarrhea, disease affecting young pheasants | 390 | 36–37 |
| Diet, meat, reduction of expense | 391 | 16–21 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| Diet, place of bread | 389 | 45–46 |
| Dietrich, W., report on experiments in pig feeding in Illinois | 388 | 25–28 |
| Dionin, use as habit-forming agent | 393 | 3 |
| Dip, arsenical, use in destruction of cattle ticks, method and formula for preparation | 378 | 22–24 |
| Dipping, value in preservation of posts, methods | 387 | 7 |
| Dips, arsenical, manufacture and use, precautions | 378 | 24 |
| preparation and use in destruction of Texas-fever ticks | 378 | 21–27 |
| Disease, loco-weed | 380 | 1–16 |
| Diseases, hog, confusion with hog cholera | 379 | 15–17 |
| nose and lungs, use of habit-forming drugs in treatment | 393 | 8–14 |
| pheasants, control methods, etc | 390 | 35–40 |
| Disinfectant, cresol solution, use in hog cholera prevention | 379 | 18 |
| Disinfectants, use in control of pheasant diseases | 390 | 35–36 |
| Disinfection, control method for pheasant diseases | 390 | 35–36 |
| Ditches, field, for irrigation, making, implements, etc | 399 | 7–9 |
| *Dolichos biflorus*, drought resistance | 388 | 11 |
| *lablab*, drought resistance | 388 | 11 |
| Dorset, M., bulletin on "Hog cholera" | 379 | 1–23 |
| Dough, preparation for bread, methods | 389 | 22–28 |
| Drill, disk, kind, cost, etc | 388 | 15 |
| use and value in oat production | 388 | 14–16 |
| Drilling, corn, study of method | 400 | 5, 6 |
| Drinks, medicated, soft, habit-forming agents | 393 | 6–8 |
| soft, cocaine and caffein content | 393 | 7–8 |
| dangerous ingredients | 393 | 6–8 |
| habit-forming agents | 393 | 6–8 |
| Drippings, savory, preparation | 391 | 20 |
| Drought resistance, dry-land agriculture, principles governing | 388 | 10–11 |
| Drug addiction, control methods, recommendation | 393 | 18–19 |
| treatment | 393 | 3 |
| addicts, total estimated number in United States | 393 | 3 |
| products, misbranding investigations, reasons | 377 | 5 |
| Drug-addiction treatments, harmful nature, notes | 393 | 15–18 |
| Drug-habit cures, responsibility of physicians in perpetuation | 393 | 18 |
| Drugs, habit-forming agents | 393 | 1–19 |
| labeling requirements | 393 | 4 |
| menace of sale and use | 393 | 1–19 |
| products containing, studies | 393 | 4–18 |
| use in treatment of nose and lung diseases | 393 | 8–14 |
| value of proper labels | 393 | 4 |
| harmful effects, investigation, method | 377 | 5 |
| results, summary | 377 | 6 |
| unfavorable symptoms produced, description | 377 | 4 |
| Dry farming, definition | 388 | 6 |
| importance of supplementing with irrigation | 388 | 11–12 |
| principles, studies | 388 | 6–12 |
| region, adaptability of various crops | 388 | 10–11 |
| summer tillage, results and value | 388 | 7–10 |
| Dry-land farming. *See* Dry farming. | | |
| Dumplings, preparation with meat stew | 391 | 21–22 |
| Durum wheat, value for bread making compared with other wheats | 389 | 37 |
| Dynamite, use in removal of stumps | 381 | 8 |
| Education, rural, relation of club work | 385 | 11–12 |
| school-extension work, exhibits, etc | 385 | 11–12 |
| Eelworms, destruction, experiments in Massachusetts | 388 | 17–18 |
| garden and greenhouse soils, destruction methods | 388 | 17–18 |
| Egyptian bees, note | 397 | 13 |
| Emulsion, crude petroleum, method and formula for preparation | 378 | 21–22 |
| England, early crossing experiments with pheasants | 390 | 12–13 |
| Enteritis, disease affecting adult pheasants | 390 | 38 |
| Entire-wheat flour and bread, composition, etc | 389 | 40–42 |
| Epilepsy remedies, harmful nature, notes | 393 | 15 |
| Epsom salts, loco-poisoning antidote | 380 | 13 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| Equipment, special, use in Camembert cheese making | 384 | 24–26, 27–28 |
| Experiment Station Work, LIV | 381 | 1–32 |
| LV | 384 | 1–32 |
| LVI | 388 | 1–32 |
| Exports, game, regulations, by States and Provinces | 376 | 45–52 |
| Factories, beet-sugar, number and capacities, by States, 1908 | 392 | 8 |
| Fallowing, summer, and continuous cropping, studies and results in Montana | 388 | 8 |
| Farm animals, locoed, treatment, experiments in Colorado | 380 | 13–14 |
| post-mortem appearance | 380 | 12 |
| *See also* Live stock. | | |
| fertilizer practice, studies in the South Atlantic States | 398 | 11–17 |
| Farming, dry, importance of supplementing with irrigation | 388 | 11–12 |
| *See* Dry farming. | | |
| muskrat, returns, possibilities, etc | 396 | 31–33 |
| Fasteners, wire, for poured cement fence posts, description, etc | 384 | 32 |
| Feeding bees, directions | 397 | 23–24 |
| calves, experiments and results | 381 | 14–23 |
| pigs, experiments in Illinois | 388 | 25–28 |
| stuff, alfalfa meal, studies of value, etc | 384 | 12–14 |
| Fence posts, cement, manufacture, methods, etc | 384 | 29–32 |
| concrete, manufacture, methods, etc | 384 | 29–32 |
| poured, comparison with tamped posts | 384 | 31–32 |
| use of reinforcements | 384 | 32 |
| tamped, comparison with poured posts | 384 | 31–32 |
| Fermentation, lessening danger of cotton-seed meal as swine food | 384 | 10–12 |
| yeast and other leavening agencies | 389 | 17–21 |
| Fertilizer constituents, various crops and soils, recommendations | 398 | 16 |
| mixtures, incompatibles, study, diagram | 388 | 5 |
| use in corn fertilizing, quantity, composition, etc | 398 | 19–21 |
| cotton fertilizing, quantity, composition, etc | 398 | 17–19 |
| oat fertilizing, quantity, composition etc. | 398 | 21–22 |
| wheat fertilizing, quantity, composition, etc | 398 | 22–23 |
| use in fertilizing cowpeas, kinds, quantity, etc | 398 | 23–24 |
| South Atlantic States, kind and quantity for various crops, etc | 398 | 12–17 |
| Fertilizers, combinations, safe, list and diagram | 388 | 5 |
| commercial, factors influencing use | 398 | 6–11 |
| use in South Atlantic States, farm practice | 398 | 1–24 |
| sugar-beet growing, management | 392 | 39–40 |
| Fertilizing corn, studies, recommendations, etc | 398 | 19–21 |
| cotton, studies, recommendations, etc | 398 | 17–19 |
| cowpeas, studies | 398 | 23–24 |
| material, availability, kind, quantity, etc | 398 | 17–24 |
| materials, indiscriminate mixing, danger, studies | 388 | 5 |
| oats, studies, recommendation, etc | 398 | 21–22 |
| rotation practices in South Atlantic States, investigations | 398 | 11 |
| wheat, studies, suggestions, etc | 398 | 22–23 |
| Fescue, meadow. *See* Meadow fescue. | | |
| *Fiber zibethicus*. *See* Muskrat. | | |
| Flavor of meat | 391 | 12–13 |
| developing and improving | 391 | 35–40 |
| Flaxseed, use in artificial feed for calves in Germany | 381 | 22 |
| Float, land-leveling, description | 392 | 12–13 |
| Flooding, irrigation of grain, methods, etc | 399 | 4–14 |
| Flour, chemical composition | 389 | 36–37 |
| gluten content, relation to quality of bread | 389 | 31–33 |
| impurities, prevention | 389 | 13–14 |
| rye, uses, value, etc | 389 | 13–14 |
| wheat, tests for quality | 389 | 13 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| Flours, bread making, value of different kinds | 389 | 13–16 |
| description, composition, etc | 389 | 40–42 |
| use in bread making | 389 | 8–16 |
| Food, use of muskrat, preparation, etc | 396 | 21–24 |
| value of meat | 391 | 5–8 |
| Forcemeat, recipe | 391 | 27 |
| Fowler's solution, loco poisoning antidote | 380 | 13, 14 |
| Fowls, jungle, ancestors of domestic chickens | 390 | 10–11 |
| origin, description, etc | 390 | 10–11 |
| France, artificial feed for calves, experiments and results | 381 | 23 |
| FULLER, P. E., bulletin on "The use of windmills in irrigation in the semiarid West" | 394 | 1–44 |
| Fungi, responsibility for decay in farm timbers | 387 | 5–6 |
| Fur supply, future, suggestions | 396 | 9 |
| Furrow irrigation, management | 392 | 14–22 |
| method, etc | 399 | 12–14 |
| Furs, dressing | 396 | 29–31 |
| use, growth, remarks | 396 | 7 |
| *Galleria mellonella*. *See* Wax moth, larger. | | |
| *Gallus*, pheasant variety | 390 | 10–11 |
| Game, bag limits, changes in laws | 376 | 9 |
| bills, Federal and State, failing of enactment in 1909 | 376 | 15–17 |
| export regulations, States and Provinces | 376 | 45–52 |
| restrictions | 376 | 8 |
| exportation, Canadian laws regulating | 376 | 37 |
| State laws prohibiting | 376 | 33–37 |
| killing, limits, list of States and Provinces | 376 | 41–44 |
| laws, enactment in 1909, provisions | 376 | 11–15 |
| for 1909 | 376 | 1–56 |
| States and Provinces, enactment in 1909, list | 376 | 6, 11–15 |
| legislation, 1909, discussion | 376 | 6–11 |
| licenses, amounts, various States and Provinces | 376 | 9–10 |
| for hunting and shipping, States and Provinces | 376 | 45–52 |
| States and Provinces requiring | 376 | 9–10 |
| open seasons, county laws for Maryland | 376 | 53–54 |
| North Carolina | 376 | 55–56 |
| notes | 376 | 53 |
| United States and Canada, 1909, table | 376 | 19–32 |
| preserves, establishment, 1909 | 376 | 10–11 |
| private, experiments in pheasant propagation | 390 | 17 |
| protection, open seasons, changes | 376 | 8 |
| refuges, establishment, 1909 | 376 | 10–11 |
| sale in close seasons, notes | 376 | 38 |
| open season | 376 | 38 |
| prohibition for entire year, list of States and Provinces | 367 | 38–41 |
| regulations | 376 | 37–41 |
| restrictions | 376 | 8–9 |
| shipment, Federal and State laws regulating | 376 | 32–37 |
| laws affecting | 376 | 32 |
| shipping, license requirements, States and Provinces | 376 | 45–52 |
| wardens, change in service | 376 | 10 |
| increase of salaries | 376 | 10 |
| Gapes, disease affecting young pheasants | 390 | 37 |
| Gasoline, use in heating brooders in New York | 381 | 23–30 |
| *Gennæus nycthemerus*, pheasant variety | 390 | 9 |
| Georgia, boys' and girls' agricultural clubs, work and results | 385 | 10 |
| Germany, artificial feeds for calves, experiments and results | 381 | 22 |
| Goldthwaite, Miss N. E., studies of jelly-making fruit juices | 388 | 30–32 |
| Goulash, Hungarian, recipe | 391 | 30 |
| Grading, comb honey, rules | 397 | 35–36 |
| Graham bread, composition, etc | 389 | 40–42 |
| flour, composition, etc | 389 | 40–42 |
| Grain, alfalfa meal as substitute, for live stock | 384 | 12–14 |
| growing under irrigation, cost and returns | 399 | 21–23 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| Grain, irrigation methods, etc | 399 | 1–23 |
| land, irrigated, value | 399 | 21 |
| Grains, use in bread making, treatment, etc | 389 | 8–16 |
| Gravy, meat, suggestions for making | 391 | 39–40 |
| GRAYBILL, H. W., bulletin on "Methods of exterminating the Texas-fever tick" | 378 | 1–30 |
| GRUBB, E. H., bulletin on "Potato culture on irrigated farms of the West" | 386 | 1–13 |
| Guava juice, study and experiments in jelly-making | 388 | 29 |
| Habit-forming agents | 393 | 1–19 |
| importation and use | 393 | 3 |
| HARTLEY, C. P., bulletin on "A more profitable corn-planting method" | 400 | 1–14 |
| Harvesting sugar beets, directions, methods, etc | 392 | 33–37 |
| Hash, casserole or Italian, recipe | 391 | 31 |
| Hay, feeding value for milch cows, comparison with mangels | 384 | 15 |
| timothy, protein content, comparison with alfalfa meal | 384 | 13–14 |
| Headache mixtures, harmfulness | 377 | 1–16 |
| Heart, stuffed, recipe | 391 | 29 |
| Herbs, flavoring, use in cooking meats, etc | 391 | 37–40 |
| Herds, dairy. *See* Dairy herds. | | |
| Heroin, use as habit-forming agent | 393 | 3 |
| in soothing sirups | 393 | 5 |
| HILLMAN, F. H., bulletin on "The adulteration of forage-plant seeds" | 382 | 1–23 |
| Hive stands, construction | 397 | 10 |
| tools, illustrations | 397 | 10–11 |
| Hog cholera | 379 | 1–23 |
| appearance of hog carcasses after death | 379 | 9–15 |
| carcasses, danger to man in examination | 379 | 9 |
| causes | 379 | 6 |
| confusion with other hog diseases | 379 | 15–17 |
| first outbreak in United States, distribution, description, damage, etc | 379 | 5 |
| introduction of germ, various methods | 379 | 6–8 |
| prevention and cure, medicine formula | 379 | 19 |
| treatment | 379 | 18–23 |
| disinfection | 379 | 18 |
| inoculation, methods and results | 379 | 19–22 |
| necessity of cooperation with State authorities | 379 | 23 |
| quarantine methods | 379 | 18 |
| symptoms, diagnosis | 379 | 8–9, 15 |
| disease, anthrax, confusion with hog cholera | 379 | 17 |
| lung worms, confusion with hog cholera | 379 | 17 |
| swill-fed hogs, responsibility of soap slops | 379 | 15–16 |
| swine plague, confusion with hog cholera | 379 | 16 |
| tuberculosis, confusion with hog cholera | 379 | 16–17 |
| diseases, similarity to hog cholera | 379 | 15–17 |
| Hogs, feeding fermented cotton-seed meal, danger | 384 | 9–11 |
| swill-feeding, danger from soap in slops | 379 | 15–16 |
| "Holyland" bees, note | 397 | 13 |
| Home, economical use of meat in | 391 | 43 |
| Honey, adulteration, laws affecting, note | 397 | 42 |
| comb, production for market, method | 397 | 32–36 |
| extracted, production, method | 397 | 31–33 |
| production, marketing, etc | 397 | 5–44 |
| notes | 397 | 30–31 |
| Honeycomb, structure, uses in bee colony, etc | 397 | 14 |
| "Honeydew" honey, source, value, etc | 397 | 31 |
| Horses, farm work, rations for winter, experiments in Michigan | 384 | 11–12 |
| wintering experiments | 384 | 11–12 |
| loco poisoning, symptoms | 380 | 11 |
| poisoning by loco weed | 380 | 9–11 |
| HOWE, F. W., bulletin on "Boys' and girls' agricultural clubs" | 385 | 1–23 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| HUNT, CAROLINE L., and C. F. LANGWORTHY, bulletin on "Economical use of meat in the home" | 391 | 1–43 |
| Hunting, licenses, requirements, States and Provinces | 376 | 45–52 |
| Idaho, dairy herds, production, profits, etc., studies | 384 | 16 |
| tests of Sixty-day oats | 395 | 24–25 |
| Illinois, annual oat production | 388 | 12 |
| boys' and girls' agricultural clubs, work and results | 385 | 9–10 |
| calves, cost of raising | 381 | 17–18 |
| experiments in pheasant propagation | 390 | 16 |
| oat production, annual increase, possibility | 388 | 15 |
| loss from smut | 388 | 13 |
| methods, yield, and value | 388 | 12, 13, 14–15 |
| tests of Sixty-day and Kherson oats | 395 | 16–17 |
| Implements for preparing land for irrigation | 399 | 7, 8–9, 11 |
| sugar-beet culture, illustrations, etc | 392 | 15, 17–18, 32, 34 |
| Indiana, experiments in pheasant propagation | 390 | 19 |
| tests of Kherson and Sixty-day oats | 395 | 14–15 |
| Inoculation, hog cholera prevention, methods and results | 379 | 19–22 |
| mold spores, method in Camembert cheese making | 384 | 27 |
| Intestinal worms, disease affecting adult pheasants | 390 | 39 |
| Iowa, oat production, methods, yield, and value | 388 | 12, 13, 15 |
| tests of Kherson and Sixty-day oats | 395 | 18 |
| Irrigated sections, beet-sugar factories | 392 | 8 |
| Irrigation, danger | 399 | 20–21 |
| implements for preparation of land for | 399 | 7, 8–9, 11 |
| importance in supplementing dry farming | 388 | 11–12 |
| grain, methods, etc | 399 | 1–23 |
| time, water requirements, etc | 399 | 14–20 |
| of sugar beets | 392 | 1–52 |
| potato growing in West, methods, etc | 386 | 8 |
| preparation of land | 392 | 11–12 |
| small areas, study of methods | 388 | 12 |
| sugar-beet growing, methods, time, etc | 392 | 12–28 |
| results of experiments | 392 | 45–47 |
| use in onion culture in Southwest | 384 | 6–7 |
| of windmills in semiarid West | 394 | 1–44 |
| water supply, use, etc | 392 | 41–43 |
| Italian bees, value, remarks | 397 | 13 |
| Italy, artificial feeds for calves, experiments and results | 381 | 20–21 |
| *Ithaginis*, pheasant variety | 390 | 10 |
| Jelly, guava, directions for making | 388 | 29 |
| making, experiments and results | 388 | 29–32 |
| with various acids | 388 | 31–32 |
| with fruit juices, studies and experiments | 388 | 30–32 |
| Juices, fruit, effects of sugar and temperature on, studies | 388 | 30 |
| Kansas, experiments in pheasant propagation | 390 | 17 |
| oat production, methods, yield, etc | 388 | 13–14 |
| tests of Sixty-day and Kherson oats | 395 | 22–23 |
| western, windmill irrigation | 394 | 37–39 |
| KEBLER, L. F., bulletin on "Habit-forming agents" | 393 | 1–19 |
| F. P. MORGAN, and PHILIP RUPP, bulletin on "Harmfulness of headache mixtures" | 377 | 1–16 |
| Kentucky bluegrass seed, adulteration, sample tests, etc | 382 | 12–14 |
| tests of Kherson and Sixty-day oats | 395 | 15 |
| Kherson oats, with sixty-day oats | 395 | 1–27 |
| Koca Nola, harmful character, note | 393 | 8 |
| Koke, harmful character, note | 393 | 8 |
| Pillsbury's, harmful character, note | 393 | 8 |
| Kola nut, caffein source | 393 | 7 |
| Kola-Ade, harmful character, note | 393 | 8 |
| Kos-Kola, harmful character, note | 393 | 8 |
| Kulthi, drought resistance | 388 | 11 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| Labeling, value for habit-forming drugs | 393 | 4 |
| Labels, misleading, for habit-forming agents | 393 | 4–5 |
| Lactic acid, use in whipping cream | 384 | 21 |
| manufacture of skim-milk buttermilk | 384 | 18–19 |
| Land, logged-off, clearing, methods and cost | 381 | 5–9 |
| Langstroth, inventor of bee hive, note | 397 | 9 |
| Langworthy, C. F., and Caroline L. Hunt, bulletin on "Economical use of meat in the home" | 391 | 1–43 |
| Lantz, David E., bulletin on "The muskrat" | 396 | 1–38 |
| *Laphophorus refulgens*, pheasant variety | 390 | 10 |
| Laws affecting bee keeping | 397 | 42 |
| game, enactment in 1909, list of States and Provinces | 376 | 6, 11–15 |
| *See also* Legislation. | | |
| League, Farmer Boys' and Girls', Texas, organization and work | 385 | 10 |
| Leaven, use in raising bread, method | 389 | 25–26 |
| Leavening agencies | 389 | 17–21 |
| Legislation, game. *See* Game laws. | | |
| protection of muskrats | 396 | 35–37 |
| *See also* Laws. | | |
| Legumes, growing, influence on use of commercial fertilizers | 398 | 6–7 |
| use as foundation crop in potato culture | 386 | 5 |
| Levelers, land, description, illustration, etc | 392 | 17–18 |
| Licenses, game, hunting and shipping, States and Provinces, requirements | 376 | 45–52 |
| hunting and shipping game, requirements, States and Provinces | 376 | 45–52 |
| Lime, use as fertilizer in South Atlantic States, notes | 398 | 10–11 |
| Lindsey, J. B., report on animal feeds for farm stock | 388 | 24–25 |
| Live stock, alfalfa meal as substitute for grain | 384 | 12–14 |
| animal feeds | 388 | 24–25 |
| oleander poisoning, experiments and results | 384 | 8–9 |
| symptoms | 384 | 9 |
| Loco disease, horses, cattle, and sheep, presence in Great Plains region | 380 | 5 |
| origin and meaning of word | 380 | 5 |
| plant, botanical classification | 380 | 5–9 |
| poisoning, post-mortem appearance of animals | 380 | 12 |
| weed, name given to poisonous plants of Leguminosæ family | 380 | 5 |
| pink, description | 380 | 8 |
| poisoning, various animals | 380 | 9–11 |
| purple, distribution, description, etc | 380 | 6–7, 8 |
| poisonous nature, studies by Plant Industry Bureau, and results | 380 | 9–11 |
| white, description and distribution | 380 | 7–8 |
| poisonous nature, studies by Plant Industry Bureau, and results | 380 | 9–11 |
| weeds, destruction by insects in various States | 380 | 15–16 |
| methods, experiments, etc | 380 | 14–16 |
| laboratory investigations, and results | 380 | 12–13 |
| Loco-weed disease | 380 | 1–16 |
| "Long-tongued bees," note | 397 | 13 |
| Lophura, pheasant variety | 390 | 10 |
| Lung diseases, treatment, use of harmful drugs | 393 | 8–14 |
| Mangels, cost, comparison with other feeding stuffs | 384 | 15 |
| feeding to dairy cows, milk yield, quality, etc | 384 | 14–15 |
| value for milch cows, comparison with hay, grain, silage, etc | 384 | 14–15 |
| Manure, barnyard, influence on use of commercial fertilizers | 398 | 8 |
| green, influence on use of commercial fertilizers | 398 | 7–8 |
| poultry, preservation and value | 384 | 5–6 |
| sugar-beet growing, use | 392 | 39–40 |
| *Margaropus annulatus*. *See* Tick, Texas-fever. | | |
| Marketing potatoes, study of methods and results | 386 | 10 |
| Marrow, preparation for the table | 391 | 20 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| MARSH, C. DWIGHT, bulletin on "The loco-weed disease" | 380 | 1–16 |
| lands, use for muskrat farming | 396 | 32–33 |
| Maryland, acclimatization of pheasants, early experiments | 390 | 13–14 |
| game, open seasons, county laws | 376 | 53–54 |
| tests of Sixty-day oats | 395 | 13 |
| Massachusetts, alfalfa meal *vs.* wheat bran as cow feed, experiments | 384 | 14 |
| calves, cost of raising | 381 | 17 |
| experiments in acclimatization of pheasants | 390 | 14–15, 18–19 |
| MCLAUGHLIN, WALTER W., bulletin on "Irrigation of grain" | 399 | 1–23 |
| McWethy, L. B., statement on Kherson oat tests | 395 | 24 |
| Meadow fescue seed, adulteration | 382 | 12 |
| use as adulterant in orchard grass seed | 382 | 11–12 |
| Meal, alfalfa, use and value as substitute for grain | 384 | 12–14 |
| cotton-seed, fermented, feeding to hogs, danger, experiments, etc | 384 | 9–11 |
| fermentation as means of lessening danger as swine food | 384 | 10–12 |
| injurious effects on hogs, studies and experiments | 384 | 9–11 |
| Meat, cheap cuts, utilization | 391 | 28–35 |
| composition of different cuts | 391 | 6–8 |
| cooking, changes, etc | 391 | 13–16 |
| different cuts, cost, etc | 391 | 8–11 |
| economical use in the home | 391 | 43 |
| flavor, developing and improving | 391 | 35–40 |
| extension in food | 391 | 21 |
| industry, by-products, use as stock foods | 388 | 24–25 |
| preparation for table, methods | 391 | 13–16 |
| prices, relation of different cuts | 391 | 8–10 |
| texture and flavor | 391 | 12–13 |
| tough, cooking, recipes | 391 | 28–35 |
| value as food | 391 | 5–8 |
| various dishes, preparation | 391 | 22–27 |
| Meats, chopped, preparation, recipes | 391 | 32–35 |
| pounded, preparation, recipes | 391 | 31–32 |
| Medicines, introduction of dangerous drugs for | 377 | 3 |
| Michigan, tests of Sixty-day and Kherson oats | 395 | 15 |
| winter rations for farm work horses, experiments and results | 384 | 11–12 |
| *Micrococcus prodigrosus*, effect on bread | 389 | 35 |
| Mildew, downy, damage to onion crop in Southwest | 384 | 8 |
| Milk, butter fat, substitutes, experiments and results | 381 | 14–23 |
| skim, substitutes, study and experiments in New York | 381 | 19 |
| use in manufacture of buttermilk | 384 | 18–19 |
| of molasses as supplement, experiment in Louisiana, and results | 381 | 14–15 |
| various grains as supplement, experiments in Virginia, and results | 381 | 15–17 |
| Milling grain, processes, products, etc | 389 | 11–13 |
| Minnesota, experiments in pheasant propagation | 390 | 17 |
| tests of Sixty-day and Kherson oats | 395 | 17–18 |
| Mite, air-sac, disease affecting adult pheasants | 390 | 39 |
| Mock duck, recipe | 391 | 27 |
| Molasses, low-grade sugar-cane, use as supplement to skim milk, experiment in Louisiana, and result | 381 | 14–15 |
| Monauls, origin, description, etc | 390 | 10, 20 |
| Montana, adaptability of various crops | 388 | 10–11 |
| tests of Sixty-day and Kherson oats | 395 | 23–24 |
| Mooers, C. A., statement on Kherson and Sixty-day oat tests in Tennessee | 395 | 16 |
| Moore, Professor, statement on Kherson and Sixty-day oat tests in Wisconsin | 395 | 17 |
| MORGAN, F. P., L. F. KEBLER, and PHILIP RUPP, bulletin on "Harmfulness of headache mixtures" | 377 | 1–16 |
| Morphine, use in soothing sirups | 393 | 4–5 |
| MORSE, GEORGE BYRON, article on "Diseases of pheasants" | 390 | 35–40 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| Moths, wax, enemies to bees, injuries, etc | 397 | 40 |
| Muskrat, classification and distribution | 396 | 10 |
| description, habits, etc | 396 | 11–21 |
| economic uses | 396 | 21–31 |
| enemies | 396 | 35 |
| farming, returns, possibilities, etc | 396 | 31–33 |
| habits, economic uses, etc | 396 | 1–38 |
| pelts, demand, sales, etc | 396 | 9 |
| sales, numbers, etc | 396 | 9, 24–26 |
| protective laws, closed seasons, etc | 396 | 35–37 |
| use as food, preparation, etc | 396 | 21–24 |
| Muskrats, destruction, protection from, etc., methods | 396 | 34–35 |
| trapping, bait traps, etc | 396 | 26–28 |
| Musquash. *See* Muskrat. | | |
| Nebraska, adaptability of various crops | 388 | 11 |
| boys' and girls' agricultural clubs, work and results | 385 | 8–9 |
| experiments with Kherson oats | 395 | 8 |
| tests of Kherson and Sixty-day oats | 395 | 22 |
| western, windmill irrigation | 394 | 37–39 |
| Nematodes. *See* Eelworms. | | |
| New Hampshire, experiments in pheasant propagation | 390 | 17 |
| Jersey, early experiments in acclimatization of pheasants | 390 | 13, 16–17 |
| Mexico, early onion raising, experiments and results | 384 | 6, 8 |
| tests of Kherson oats | 395 | 23 |
| York, boys' and girls' agricultural clubs, work and results | 385 | 7–8 |
| experiments in acclimatization of pheasants | 390 | 15 |
| skim-milk substitutes, experiments and results | 381 | 19 |
| tests of Sixty-day oats | 395 | 14 |
| use of gasoline-heated colony brooders | 381 | 23–30 |
| North Carolina, experiments in use of fermented cotton-seed meal as swine food | 384 | 10–11 |
| game, open seasons, county laws | 376 | 55–56 |
| use of rolled oats for milk in calf feeding | 381 | 18, 19 |
| Dakota, tests of Sixty-day and Kherson oats | 395 | 18–20 |
| Nose diseases, treatment, use of harmful drugs | 305 | 8–14 |
| Oat flour, use in bread making, note | 389 | 14 |
| production, use and value of drill in seeding | 388 | 14–16 |
| seeding, comparison with disk drill with broadcast seeder | 388 | 14–16 |
| yields, comparison of Sixty-day and Kherson oats | 395 | 26–27 |
| methods for increasing | 388 | 13–16 |
| Oats, annual production in Illinois | 388 | 12 |
| drilling, comparison with broadcasting | 388 | 14–16 |
| early, adaptability to corn belt | 395 | 7–8 |
| fertilizing, studies and suggestions | 398 | 21–22 |
| Illinois, yield per acre, and value | 388 | 12, 13, 14–15 |
| Iowa, yield per acre, and value | 388 | 12, 13, 15 |
| Kansas, yield per acre, etc | 388 | 13–14 |
| Kherson, adaptability to various States | 395 | 12–13 |
| amount of seed per acre | 395 | 12 |
| comparison with other oats | 395 | 9–12 |
| description, characteristics, etc | 395 | 9–12 |
| experiments in various States | 395 | 8, 13–26 |
| introduction into United States | 395 | 8 |
| origin, history, and distribution | 395 | 8–9 |
| value for nurse crops | 395 | 12, 16 |
| methods of seeding in various States | 388 | 13–16 |
| production in Canada, methods, yield, etc | 388 | 13, 14, 15 |
| rolled, substitute for milk in calf feeding | 381 | 18–19 |
| Sixty-day, adaptability to various States | 395 | 12–13 |
| amount of seed per acre | 395 | 12 |
| and Kherson | 395 | 1–27 |
| comparison of yields | 395 | 26–27 |
| chemical composition, comparison with other oats | 395 | 10–12 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| Oats, Sixty-day, comparison with other oats | 395 | 9–12 |
| description, characteristics, etc | 395 | 9–12 |
| experiments in various States | 395 | 13–26 |
| introduction into United States | 395 | 8–9 |
| origin, history, and distribution | 395 | 8–9 |
| similarity to Kherson oats | 395 | 8 |
| tillage methods, experiments in Montana | 388 | 8 |
| value as nurse crop | 395 | 12, 16 |
| yield in corn belt, influences controlling | 395 | 7–8 |
| Ogilvie-Grant, statement on purity of pheasant stock | 390 | 13 |
| Ohio, experiments in acclimatization of pheasants | 390 | 15 |
| tests of Sixty-day oats | 395 | 14 |
| Oil, cod-liver, use as substitute for butter fat in feeding calves | 381 | 19–20 |
| Oklahoma, tests of Kherson and Sixty-day oats | 395 | 23 |
| OLDYS, HENRY, bulletin on "Pheasant raising in the United States" | 390 | 1–34 |
| T. S. PALMER, and C. E. BREWSTER, bulletin on "Game laws for 1909" | 376 | 1–56 |
| Oleander, growing, danger to human beings and live stock | 384 | 8–9 |
| poisoning farm animals, experiments and results | 384 | 9 |
| symptoms | 384 | 9 |
| of live stock, experiments and results | 384 | 8–9 |
| Oliver, Robert, early importation of pheasants | 390 | 13–14 |
| Onion growing, adaptability of soil in Southwest | 384 | 6–7 |
| Southwest, methods | 384 | 6–7 |
| Onions, early, growing in Southwest, varieties, yield, etc | 384 | 6–8 |
| Northwest, forcing methods | 384 | 8 |
| Southwest, damage by mildew and thrips | 384 | 8 |
| Ontario, use of cocoa shells for skim milk, experiments | 381 | 19 |
| Open seasons, game protection changes | 376 | 8 |
| Opium, importation into United States | 393 | 3 |
| use as habit-forming agent | 393 | 3 |
| in soothing sirups | 393 | 5 |
| Orchard grass seed, adulteration and adulterants | 382 | 11–12 |
| Oregon, pheasant raising, experiments in acclimatization | 390 | 14, 17, 18, 20 |
| tests of Sixty-day oats | 395 | 26 |
| Painting, value in preservation of posts, methods | 387 | 7 |
| PALMER, T. S., HENRY OLDYS, and C. E. BREWSTER, bulletin on "Game laws for 1909" | 376 | 1–56 |
| Pasting, disease affecting young pheasants | 390 | 36 |
| Pelts, muskrat, demand, sales, etc | 396 | 9, 24–26 |
| preparation for market | 396 | 29–31 |
| Pennsylvania, alfalfa meal, comparison with timothy hay and wheat bran, experiments and results | 384 | 13–14 |
| experiments in pheasant propagation | 390 | 17 |
| shelter-grown tobacco, production, experiments, yield cost, etc | 381 | 9–11 |
| tests of Sixty-day and Kherson oats | 395 | 13–14 |
| Petroleum, crude, use in destruction of Texas-fever ticks | 378 | 21 |
| *Phaseolus aconitifolius*, value for hay production | 388 | 11 |
| *Phasianus colchicus*, pheasant variety | 390 | 5, 6, 11–13 |
| × *torquatus*, pheasant variety | 390 | 5–6 |
| *hagenbecki*, pheasant variety | 390 | 19 |
| *mongolicus*, pheasant variety | 390 | 7 |
| *principalis*, pheasant variety | 390 | 19 |
| *torquatus*, pheasant variety | 390 | 5, 6, 11–13 |
| *versicolor*, pheasant variety | 390 | 6–7, 12–13 |
| Pheasant, Argus, origin, description, value, etc | 390 | 10 |
| blood, origin | | 10 |
| Bohemia, description | 390 | 8 |
| China. *See* Pheasant, ringneck. | | |
| *torquatus*. *See* Pheasant, ringneck. | | |
| Chinese. *See* Pheasant, ringneck. | | |
| ringneck. *See* Pheasant, ringneck. | | |
| dark-necked. *See* Pheasant, English. | | |
| Denny. *See* Pheasant, ringneck. | | |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| Pheasant, eared, origin, description, value, etc | 390 | 7, 8, 9–10 |
| eggs, shipment methods | 390 | 34 |
| English, origin, early history, distribution, etc | 390 | 5, 6, 11–13 |
| ringneck, origin, description, etc | 390 | 5–6 |
| firebacks, origin, description, etc | 390 | 10 |
| golden, origin, description, etc | 390 | 8–9, 20 |
| Hagenbeck, description | 390 | 19 |
| horned, origin, description, etc | 390 | 9, 10 |
| Hungarian. *See* Pheasant, English. | | |
| kalijes. *See* Pheasant, silver. | | |
| koklass, origin | | 10 |
| Lady Amherst, origin, description, etc | 390 | 8–9, 20 |
| Manchurian eared, description | 390 | 19 |
| *See also* Pheasant, eared. | | |
| monauls, origin, description, etc | 390 | 10, 20 |
| Mongolian, origin, description, etc | 390 | 5, 7 |
| value in game preserves | 390 | 19 |
| *See also* Pheasant, ringneck; Pheasant, English ringneck. | | |
| Oregon. *See* Pheasant, ringneck. | | |
| peacock, origin | 390 | 10 |
| *phasianus colchicus*, origin, early history, distribution, etc | 390 | 11–12 |
| Prince of Wales, introduction into England and United States | 390 | 19 |
| propagation, earliest history | 390 | 12 |
| pens, description | 390 | 21–23 |
| raising in Europe, history | 390 | 11–13 |
| United States | 390 | 1–40 |
| Reeves, description, distribution, etc | 390 | 8–9, 19 |
| ringneck, origin, early history, distribution, etc | 390 | 5, 6, 11–13 |
| *See also* Pheasant, English ringneck. | | |
| silver, origin, description, etc | 390 | 9, 19–20 |
| white, description | 390 | 8 |
| Pheasants, acclimatization in United States, early history and progress | 390 | 13–17 |
| adult, diseases affecting | 390 | 38–40 |
| bad habits, control methods | 390 | 24 |
| care, food, etc., in propagation | 390 | 23–25 |
| of young, methods, etc | 390 | 28–30 |
| in winter | 390 | 33 |
| diseases affecting, control methods | 390 | 35–40 |
| early crossing experiments in England | 390 | 12–13 |
| food for young | 390 | 30–33 |
| game, study, description, etc | 390 | 19–20 |
| hatching, methods, etc | 390 | 26–28 |
| hybridizing, experiments in England | 390 | 12–13 |
| in field and covert | 390 | 17–19 |
| live, shipment methods | 390 | 34 |
| marketing, shipment, etc | 390 | 33–34 |
| mating, season, methods, etc | 390 | 25–26 |
| methods of obtaining stock | 390 | 20 |
| pens for propagation, description | 390 | 21–23 |
| prices | 390 | 7, 8, 9–10, 20–21 |
| propagation, methods | 390 | 20–23 |
| protection from enemies | 390 | 23 |
| shooting methods | 390 | 19 |
| varieties, origin, habits, etc | 390 | 5–11 |
| young, diseases affecting | 390 | 36–37 |
| food | 390 | 30–33 |
| Phenacetin, character and use | 377 | 3 |
| harmful effects, investigation, results, summary | 377 | 6–7 |
| introduction as medicine | 377 | 3 |
| poisoning, number of cases, symptoms, fatalities, etc | 377 | 14–16 |
| use as habit-forming agent | 393 | 3 |
| dose, harmfulness, etc., reports of physicians | 377 | 6–9 |
| objects | 377 | 4–5 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| PHILLIPS, E. F., bulletin on "Bees" | 397 | 1–44 |
| Physicians, information regarding headache mixtures | 377 | 6–9 |
| Pig feeding, methods, experiments in Illinois | 388 | 25–28 |
| feeds, various kinds, value, etc | 388 | 25–28 |
| Pigs, carbohydrate requirements | 388 | 27 |
| protein requirements | 388 | 27 |
| water requirements | 388 | 26 |
| Planting, potatoes, methods, etc | 386 | 6–7 |
| Plants, forage, seeds and their adulterants, description | 382 | 15–21 |
| Plow, beet, description, illustration, etc | 392 | 34–35 |
| Plowing, sugar-beet growing | 392 | 28–29 |
| Plows, use in furrow irrigation, description, illustrations, etc | 392 | 15–16 |
| Pneumonia, disease affecting adult pheasants | 390 | 38 |
| Pod spot, fungous disease of bean | 388 | 21–23 |
| Poisoning, farm animals, oleander, symptoms | 384 | 9 |
| headache mixtures, number of cases, distribution, fatalities, etc | 377 | 9–16 |
| live stock, oleander | 384 | 8–9 |
| *Polyplectron*, pheasant variety | 390 | 10 |
| Posts, cold-bath treatment as preservative, methods | 387 | 9 |
| creosoting methods, cooperative experiments in various States | 387 | 5 |
| fence, creosote impregnation, methods of treatment | 387 | 9–10 |
| treating tanks, description | 387 | 10–11 |
| treatment, preparation | 387 | 11–13 |
| creosoting, cost, value, etc | 387 | 16–17 |
| prolonging life, methods | 387 | 6–17 |
| *See also* Fence posts. | | |
| Potato cellars, building, methods | 386 | 9 |
| culture, irrigated farms of West | 386 | 1–13 |
| irrigation in West, methods, etc | 386 | 8 |
| soil preparation | 386 | 5 |
| planter, type, study | 386 | 6–7 |
| planting, cut seed *vs.* whole seed, study | 386 | 7 |
| methods | 386 | 6–7 |
| use of machine, methods, etc | 386 | 6–7 |
| Potatoes, adaptability to Montana | 388 | 10 |
| cultivation, time and method | 386 | 7 |
| irrigated districts, scientific production, profits | 386 | 13 |
| marketing, study of methods and results | 386 | 10 |
| seed, comparison of whole and cut | 386 | 7 |
| selection, methods and results | 386 | 10–12 |
| storage, cellars, method of building | 386 | 9 |
| varieties, adaptability to western conditions | 386 | 12 |
| yields, study of possibilities | 386 | 12–13 |
| Poultry manure, preservation and value | 384 | 5–6 |
| use of chemicals and sawdust | 384 | 5–6 |
| Powder, use in removal of stumps from logged-off land | 381 | 8 |
| Preserves, game, private, experiments in pheasant propagation | 390 | 17 |
| Pruning, experiments in Colorado | 388 | 19–21 |
| importance and advantages of system | 388 | 19–21 |
| influence upon fruit-bearing habits of trees | 388 | 20 |
| methods | 388 | 20 |
| tools | 388 | 20–21 |
| Publications, junior agricultural clubs, list | 385 | 19–22 |
| bee-keeping | 397 | 43–44 |
| *Pucrasia*, pheasant variety | 390 | 10 |
| Pudding, Yorkshire, with beef, recipe | 391 | 26 |
| Puddling, reservoir method | 394 | 31–32 |
| Pumping water for irrigation, use of windmills | 394 | 1–44 |
| power requirements | 394 | 12–14 |
| with windmills, load regulators | 394 | 22–23 |
| Pumps, windmill, selection, etc | 394 | 25–28 |
| Punic bees, note | 397 | 13 |
| Queen bee, habits, use in colony, etc | 397 | 16–17 |
| bees, shipping, etc | 397 | 40–41 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| Rainfall, damage to soil surface from light showers | 388 | 8 |
| minimum, experiments in dry farming | 388 | 6 |
| retention in dry farming, conditions governing | 388 | 8 |
| soil retention, conditions governing | 388 | 8 |
| Rasmussen, F., discussion of farm butter making | 384 | 22, 23 |
| Rations, cows, daily, cost, etc., studies in Connecticut | 384 | 16–18 |
| horses, study and experiments in Michigan | 384 | 11–12 |
| Rattleweeds, studies | 380 | 9 |
| "Red-clover bees," note | 397 | 13 |
| Reinforcements, poured fence posts, varieties, description, etc | 384 | 32 |
| Remedies. *See* specific diseases for which required. | | |
| Reservoirs, erosion of banks, prevention | 394 | 33 |
| windmill plants, size, construction, etc | 394 | 28–35 |
| Ripening, Camembert cheese making, description of process | 384 | 28–29 |
| room, Camembert cheese manufacture, construction, etc. | 384 | 28–29 |
| Roast, casserole, recipe | 391 | 31 |
| pot, recipes | 391 | 29–30 |
| ROEDING, F. W., bulletin on "Irrigation of sugar beets" | 392 | 1–52 |
| Roots, various kinds, feeding value, comparison with mangels | 384 | 14–15 |
| Rotation, crop, approved method | 388 | 9–10 |
| study of practices in South Atlantic States | 398 | 11 |
| sugar-beet growing | 392 | 38–40 |
| Rothschild, Walter, breeder of pure Mongolian pheasants | 390 | 19 |
| Roup, disease affecting adult pheasants | 390 | 38 |
| "Run-off," cause of water loss in heavy storms | 388 | 8 |
| RUPP, PHILIP, L. F. KEBLER, and F. P. MORGAN, bulletin on "Harmfulness of headache mixtures" | 377 | 1–16 |
| Rust, fungous disease of bean | 388 | 21–23 |
| Rye flour, use in bread making, etc | 389 | 14 |
| Rye-grass, English, use as adulterant in orchard grass seed | 382 | 11–12 |
| Salads, meat, preparation | 391 | 25–26 |
| "Salt-rising" bread. *See* Bread, "salt-rising" | 389 | 21 |
| Salts, epsom. *See* Epsom salts. | | |
| Sauces, meat, recipes | 391 | 39–40 |
| Scab disease affecting adult pheasants | 390 | 39 |
| Scotch barm, use in bread making | 389 | 24–25 |
| Scraper, adjustable, for making field ditches for irrigation | 399 | 8 |
| Screenings, use as adulterant in red clover seed | 382 | 8 |
| Scurfy legs, disease affecting adult pheasants | 390 | 38–39 |
| Seed adulteration, effects | 382 | 6 |
| nature and forms | 382 | 5–6 |
| alfalfa, adulteration and adulterants | 382 | 8–9 |
| description | 382 | 17–18 |
| alsike clover, adulteration and adulterants | 382 | 10–11 |
| awnless brome-grass, description | 382 | 20 |
| brome-grass, awnless, adulteration, and adulterants | 382 | 14–15 |
| Canada bluegrass, description | 382 | 21 |
| use as adulterant in Kentucky bluegrass seed | 382 | 12–14 |
| cheat, description | 382 | 20 |
| chess, description | 382 | 20 |
| clover, alsike, description | 382 | 17 |
| bur, description | 382 | 18–19 |
| red, description | 382 | 16–17 |
| sweet, description | 382 | 19 |
| English bluegrass, description | 382 | 19 |
| rye-grass, description | 382 | 19–20 |
| use as adulterant in brome-grass | 382 | 14–15 |
| meadow fescue seed | 382 | 12 |
| Italian rye-grass, description | 382 | 20 |
| Kentucky bluegrass, adulteration and adulterants | 382 | 12–14 |
| laboratory tests | 382 | 12–14 |
| description | 382 | 20–21 |
| meadow fescue, adulteration | 382 | 12 |
| description | 382 | 19 |
| use as adulterant in brome-grass | 382 | 14–15 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| Seed, orchard grass, adulteration and adulterants | 382 | 11–12 |
| seed laboratory tests | 382 | 11–12 |
| description | 382 | 19 |
| use as adulterant in meadow fescue seed | 382 | 12 |
| potatoes, importance of careful selection | 386 | 10–12 |
| selection, methods, and results | 386 | 10–12 |
| red clover, adulteration and adulterants | 382 | 7–8 |
| redtop, adulteration | 382 | 14 |
| testing for adulterants, method | 382 | 21–23 |
| timothy, use as adulterant in redtop seed | 382 | 14 |
| trefoil, yellow, description | 382 | 18 |
| Seeding grain crops, relation to water supply | 399 | 20 |
| sugar-beet growing | 392 | 29–31 |
| Seeds, forage-plant, adulteration | 382 | 5–23 |
| kinds and adulterants | 382 | 7–15 |
| and their adulterants, description | 382 | 15–21 |
| germination, time required | 382 | 22–23 |
| testing for adulterants, laboratory locations | 382 | 23 |
| Semiarid lands, West, use of windmills in irrigation | 394 | 1–44 |
| Serum, hog cholera preventive, production by various State governments | 379 | 22–23 |
| production method | 379 | 20 |
| use in inoculation, hog cholera prevention | 379 | 19–22 |
| Sheep, loco poisoning, symptoms | 380 | 12 |
| poisoning by loco weed | 380 | 9–11 |
| Shingle preservatives, antiseptic, use, methods, apparatus, etc | 387 | 18–19 |
| nonantiseptic, application methods | 387 | 18 |
| Shingles, creosoted, coloring, methods | 387 | 19 |
| prolonging life, methods | 387 | 17–19 |
| Silage, feeding value for milch cows, comparison with mangels | 384 | 15 |
| Siloing beets, directions | 392 | 38 |
| Sirups, soothing, danger in use | 393 | 4–6 |
| habit-forming ingredients | 393 | 4–6 |
| list | 393 | 5 |
| use, control methods | 393 | 6 |
| Sixty-day oats and Kherson oats | 395 | 1–27 |
| Skim milk. *See* Milk. | | |
| Skins. *See* Pelts. | | |
| Sled, furrowing, use in irrigation, description, etc | 292 | 18 |
| Smoker, bee, construction | 397 | 10 |
| Smut, damage in oat production in Illinois | 388 | 13 |
| Snow, Miss Jenny H., studies of fruit juices and jelly making | 388 | 30 |
| Sodium cacodylate, loco poisoning antidote | 380 | 14 |
| Soil, character, influence on use of commercial fertilizers | 398 | 8–10 |
| damage to surface from light showers | 388 | 8 |
| humus content, method of restoration | 388 | 9–10 |
| preparation and cultivation methods, influence on use of fertilizers | 398 | 8–10 |
| for potato culture | 386 | 5 |
| sugar-beet growing, requirements | 392 | 9–11 |
| Soils, adaptability for onion culture in Southwest | 384 | 6–7 |
| South Atlantic States, farm practice in use of commercial fertilizers | 398 | 1–24 |
| Dakota, tests of Kherson and Sixty-day oats | 395 | 20–21 |
| Sparrows, English, destruction at nests | 383 | 7–8 |
| by baiting | 383 | 9 |
| poisoning, methods | 383 | 10–11 |
| shooting | 383 | 9–10 |
| trapping | 383 | 9 |
| methods | 383 | 1–11 |
| of fruits, flowers, etc | 383 | 5–7 |
| insects, food habits, etc | 383 | 5 |
| weed seeds | 383 | 5 |
| economic relations | 383 | 5 |
| increase, methods of prevention | 383 | 7 |
| utilization for food | 383 | 10 |
| Spices, use in cooking meats, etc | 391 | 37–40 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| Spores, mold, inoculation in Camembert cheese making, method | 384 | 27 |
| Spraying fruit trees in bloom, laws affecting, note | 397 | 42 |
| method, Texas-fever ticks | 378 | 24–27 |
| Starch, use in artificial feed for calves | 381 | 22 |
| Starter, buttermilk manufacture | 384 | 18–19 |
| Starters, use in manufacture of Camembert cheese | 384 | 26 |
| Statistics, junior agricultural clubs | 385 | 22–23 |
| Steak, tough, preparation | 391 | 33 |
| Stone, G. E., report on eelworm destruction experiments in Massachusetts | 388 | 17–18 |
| Storage, potatoes, cellars and method of building | 386 | 9 |
| Strychnine, loco poisoning antidote | 380 | 14 |
| Stuffing meats, recipe | 391 | 27 |
| Stumps, removal, cost | 381 | 6–7, 8–9 |
| from logged-off land, methods | 381 | 5–9 |
| Subirrigation, method | 392 | 24–25 |
| Sugar beets. *See* Beets. | | |
| consumption in United States, 1909 | 392 | 8 |
| Swarming, bees, definition, description, etc | 397 | 15 |
| prevention, increase, etc., management | 397 | 26–29 |
| Swine, feeding experiments in Illinois | 388 | 25–28 |
| Tankage, use as feed for dairy cows, experiments in Massachusetts | 388 | 24 |
| Tappoons for irrigation ditches | 392 | 25 |
| Tar, coal, use in preservation of farm timbers | 387 | 6, 7, 9 |
| petroleum, use in preservation of farm timbers | 387 | 6, 7, 9 |
| Taylor, F. W., introduction of Kherson oats into Nebraska | 395 | 8 |
| Tegetmeir, W. B., description of Hagenbeck pheasant | 390 | 19 |
| Ten Eyck, Prof., statement on Kherson and Sixty-day oat tests | 395 | 22–23 |
| Tennessee, dairy herds, production, profit, etc., studies | 384 | 16 |
| tests of Kherson oats | 395 | 15–16 |
| Texas, boys' and girls' agricultural clubs, work and results | 385 | 10 |
| fever tick. *See* Tick, Texas-fever. | | |
| onion growing, early, experiments and results | 384 | 6, 8 |
| tests of Sixty-day and Kherson oats | 395 | 23 |
| Thinning sugar beets | 392 | 31–32 |
| Thompson, H., clearing logged-off land, studies and data | 381 | 5, 8, 9 |
| Thrips, damage to onion crop in Southwest | 384 | 8 |
| Tick, cattle. *See* Tick, Texas-fever. | | |
| Texas-fever, destruction, dipping, spraying, etc | 378 | 19–30 |
| eradication, benefits | 378 | 5–7 |
| by pasture rotation | 378 | 12–19 |
| starvation, time required | 378 | 11 |
| methods | 378 | 10–27 |
| on uninfested fields, time required | 378 | 12 |
| time required | 378 | 11–19 |
| extermination, methods | 378 | 1–30 |
| life history, development, etc | 378 | 7–10 |
| "Tickides," oils, arsenic, etc., use in destruction of Texas-fever ticks | 378 | 19–30 |
| Tillage, dry farming, principles and methods, studies | 388 | 7–10 |
| summer, experiments in Nebraska | 388 | 8–9 |
| methods | 388 | 9–10 |
| value and results in dry farming | 388 | 7–10 |
| for fall-sown crops | 388 | 8–10 |
| Timbers, farm, preservation, methods | 387 | 5–6 |
| preservative treatment | 387 | 1–19 |
| Toast, dry, digestibility, etc | 389 | 45 |
| Tobacco, Connecticut Havana, seed improvement, experiments in Wisconsin, and results | 381 | 12–14 |
| dark, crop rotation experiments in Virginia, and results | 381 | 11–12 |
| culture, experiments in Virginia, and results | 381 | 11–12 |
| fertilization experiments in Virginia, and results | 381 | 11–12 |
| improvement work | 381 | 9–14 |
| seed, improvement, experiments in Wisconsin, and results | 381 | 12–14 |
| selection, experiments in Wisconsin, and results | 381 | 12–14 |
| testing, method | 381 | 13–14 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| Tobacco, seed-plant selection, method | 381 | 12–13 |
| Sumatra leaf, shelter-grown, production in Pennsylvania, experiments | 381 | 9–11 |
| Tobacco-habit cures, harmful nature, notes | 393 | 15 |
| Towers, windmill, selection | 394 | 23–24 |
| *Tragopan satyra*, pheasant variety | 390 | 9, 10 |
| Trapping, muskrat, bait, traps, etc | 396 | 26–28 |
| Traps, muskrat, description, construction, management, etc | 396 | 27–28 |
| Trefoil, yellow, relation to red clover and alfalfa, description | 382 | 5 |
| use as adulterant in alfalfa seed | 382 | 8 |
| red clover seed | 382 | 7 |
| Tuberculosis, disease affecting adult pheasants | 390 | 39–40 |
| Tunisian bees, note | 397 | 13 |
| University, Cornell, College of Agriculture, responsibility for growth of boys' and girls' clubs | 385 | 7 |
| Unleavened bread. *See* Bread. | | |
| Utah, experiments in pheasant propagation | 390 | 17 |
| tests of Sixty-day and Kherson oats | 395 | 25 |
| Vat, dipping, Texas-fever tick extermination, specifications and materials | 378 | 27–30 |
| "Veal birds," recipe | 391 | 27 |
| cooking, recipe | 391 | 35, 39 |
| Vegetables, flavoring, use in cooking meats, etc | 391 | 37–40 |
| Vermont, alfalfa meal *vs.* wheat bran as cow feed, experiments | 384 | 14 |
| experiments in pheasant propagation | 390 | 17 |
| Vinegar, use in cooking meats | 391 | 31 |
| Virginia, dark tobacco, culture, experiments in rotation and fertilization | 381 | 11–12 |
| grain as supplement to skim milk, experiments and results | 381 | 15–17 |
| Viscogen, use in whipping cream | 384 | 21 |
| Wadsworth, W. A., distribution of pheasants in New York | 390 | 15 |
| Walker, H. C., statement on Sixty-day oat tests | 395 | 24 |
| *Walshia amorphella*, insect enemy of loco weeds | 380 | 15–16 |
| WARBURTON, C. W., bulletin on "Sixty-day and Kherson oats" | 395 | 1–27 |
| Washington, tests of Sixty-day and Kherson oats | 395 | 25–26 |
| Water, flow through pipes, friction, table | 394 | 12–14 |
| supply, irrigation, uses, etc | 392 | 41–43 |
| semiarid West, sources, etc | 394 | 5–6 |
| Wax moth, larger, enemy to bees, injuries, etc | 397 | 40 |
| lesser, enemy to bees, injury, etc | 397 | 40 |
| Well casings, selection, etc | 394 | 7–9 |
| points, use, caution, etc | 394 | 11 |
| Wells, bored, sinking, casings, etc | 394 | 9–11 |
| dug, curbing, etc | 394 | 10 |
| sinking in semiarid West | 394 | 5–12 |
| Wheat bran, protein content, comparison with alfalfa meal, experiments | 384 | 13–14 |
| fertilizing, studies and suggestions | 398 | 22–23 |
| products of roller milling | 389 | 13 |
| rolling *vs.* harrowing, experiments in Nebraska | 388 | 16–17 |
| spring, tillage methods, experiments in Montana | 388 | 8 |
| structure, composition, milling, etc | 389 | 8–14 |
| summer tillage experiments in Nebraska | 388 | 9 |
| varieties, adaptability to Montana | 388 | 10 |
| winter, comparison of rolling with harrowing | 388 | 16–17 |
| Whipple, O. B., statement on pruning | 388 | 19–21 |
| White comb, disease affecting adult pheasants | 390 | 39 |
| H. H., statement on Sixty-day oat tests | 395 | 26 |
| Williams, C. G., statement on Sixty-day oat test in Ohio | 395 | 14 |
| WILLIS, C. P., bulletin on "The preservative treatment of farm timbers" | 387 | 1–19 |
| Wilt, fungous disease of bean | 388 | 21–23 |
| Wind velocities, Cheyenne, Wyo., 1904–1908, table | 394 | 14–15 |

| | Farmers' Bull. No. | Page. |
|---|---|---|
| Windmill, type used in California, description | 394 | 42–44 |
| Windmills, capacity, size, etc | 394 | 12–23 |
| erection, directions | 394 | 24–25 |
| maintenance | 394 | 35–36 |
| mechanical features and tests | 394 | 21–22 |
| number in use in irrigated districts | 394 | 36–44 |
| use in irrigation in semiarid West | 394 | 1–44 |
| Wintering farm work horses, experiments | 384 | 11–12 |
| Wisconsin, tests of Sixty-day and Kherson oats | 395 | 17 |
| tobacco seed improvement, seed selection, experiments and results | 381 | 12–14 |
| Wiseola, harmful character, note | 393 | 8 |
| Worms, intestinal, disease affecting adult pheasants | 390 | 39 |
| Wyoming, adaptability of various crops | 388 | 11 |
| Kherson oats, tests | 395 | 24 |
| Yeast, fermentation theory, etc | 389 | 17–21 |
| substitutes | 389 | 20–21 |

Bulle
of the
gate i
of the
those w
apply t
D. C.,
Mexico
Station
ment s

22. The
27. Fla
28. Wee
30. Gra
32. Silo
34. Mea
35. Pota
36. Cott
42. Fac
44. Com
48. The
49. She
51. Stan
52. The
54. Som
55. The
56. Exp
60. Met
61. Aspe
62. Mar
63. Care
64. Duc
65. Exp
69. Exp
73. Exp
77. The
78. Exp
79. Exp
81. Corn
82. The
83. Tob
84. Exp
85. Fish
86. Thir
87. Exp
88. Alk
91. Pota
92. Exp
93. Sug
96. Rais
97. Exp
99. Inse
100. Hog
101. Mill
103. Exp
104. Not
105. Exp
106. Bree
110. Rice
113. The
114. Exp
118. Grap
119. Exp
120. Inse
121. Bea
122. Exp
124. Exp
126. Prac
127. Imp
128. Eggs
131. Hou
ga
133. Exp
134. Tree
135. Sorg
137. The
138. Irrig
139. Emm
140. Pine
142. Prin
of
144. Exp
145. Carb
149. Exp
150. Clea
152. Scab

XUM

## FARMERS' BULLETINS.

Bulletins in this list will be sent free, so long as the supply lasts, to any resident of the United States, on application to his **Senator, Representative, or Delegate in Congress,** or to the Secretary of Agriculture, Washington, D. C. Because of the limited supply, applicants are urged to select only a few numbers, choosing those which are of special interest to them. Residents of foreign countries should apply to the Superintendent of Documents, Government Printing Office, Washington, D. C., who has these bulletins for sale. Price 5 cents each to Canada, Cuba, and Mexico; 6 cents to other foreign countries. The bulletins entitled "Experiment Station Work" give briefly the results of experiments performed by the State experiment stations.

22. The Feeding of Farm Animals.
27. Flax for Seed and Fiber.
28. Weeds: And How to Kill Them.
30. Grape Diseases on the Pacific Coast.
32. Silos and Silage.
34. Meats: Composition and Cooking.
35. Potato Culture.
36. Cotton Seed and Its Products.
42. Facts About Milk.
44. Commercial Fertilizers.
48. The Manuring of Cotton.
49. Sheep Feeding.
51. Standard Varieties of Chickens.
52. The Sugar Beet.
54. Some Common Birds.
55. The Dairy Herd.
56. Experiment Station Work—I.
60. Methods of Curing Tobacco.
61. Asparagus Culture.
62. Marketing Farm Produce.
63. Care of Milk on the Farm.
64. Ducks and Geese.
65. Experiment Station Work—II.
69. Experiment Station Work—III.
73. Experiment Station Work—IV.
77. The Liming of Soils.
78. Experiment Station Work—V.
79. Experiment Station Work—VI.
81. Corn Culture in the South.
82. The Culture of Tobacco.
83. Tobacco Soils.
84. Experiment Station Work—VII.
85. Fish as Food.
86. Thirty Poisonous Plants.
87. Experiment Station Work—VIII.
88. Alkali Lands.
91. Potato Diseases and Treatment.
92. Experiment Station Work—IX.
93. Sugar as Food.
96. Raising Sheep for Mutton.
97. Experiment Station Work—X.
99. Insect Enemies of Shade Trees.
100. Hog Raising in the South.
101. Millets.
103. Experiment Station Work—XI.
104. Notes on Frost.
105. Experiment Station Work—XII.
106. Breeds of Dairy Cattle.
110. Rice Culture in the United States.
113. The Apple and How to Grow It.
114. Experiment Station Work—XIV.
118. Grape Growing in the South.
119. Experiment Station Work—XV.
120. Insects Affecting Tobacco.
121. Beans, Peas, and Other Legumes as Food.
122. Experiment Station Work—XVI.
124. Experiment Station Work—XVII.
126. Practical Suggestions for Farm Buildings.
127. Important Insecticides.
128. Eggs and Their Uses as Food.
131. Household Tests for Detection of Oleomargarine and Renovated Butter.
133. Experiment Station Work—XVIII.
134. Tree Planting on Rural School Grounds.
135. Sorghum Sirup Manufacture.
137. The Angora Goat.
138. Irrigation in Field and Garden.
139. Emmer: A Grain for the Semiarid Regions.
140. Pineapple Growing.
142. Principles of Nutrition and Nutritive Value of Food.
144. Experiment Station Work—XIX.
145. Carbon Bisulphid as an Insecticide.
149. Experiment Station Work—XX.
150. Clearing New Land.
152. Scabies of Cattle.
154. The Home Fruit Garden: Preparation and Care.
155. How Insects Affect Health in Rural Districts.
156. The Home Vineyard.
157. The Propagation of Plants.
158. How to Build Small Irrigation Ditches.
162. Experiment Station Work—XXI.
164. Rape as a Forage Crop.
166. Cheese Making on the Farm.
167. Cassava.
169. Experiment Station Work—XXII.
170. Principles of Horse Feeding.
172. Scale Insects and Mites on Citrus Trees.
173. Primer of Forestry. Part I: The Forest.
174. Broom Corn.
175. Home Manufacture and Use of Unfermented Grape Juice.
176. Cranberry Culture.
177. Squab Raising.
178. Insects Injurious in Cranberry Culture.
179. Horseshoeing.
181. Pruning.
182. Poultry as Food.
183. Meat on the Farm: Butchering, Curing, and Keeping.
185. Beautifying the Home Grounds.
186. Experiment Station Work—XXIII.
187. Drainage of Farm Lands.
188. Weeds Used in Medicine.
190. Experiment Station Work—XXIV.
192. Barnyard Manure.
193. Experiment Station Work—XXV.
194. Alfalfa Seed.
195. Annual Flowering Plants.
196. Usefulness of the American Toad.
197. Importation of Game Birds and Eggs for Propagation.
198. Strawberries.
199. Corn Growing.
200. Turkeys.
201. Cream Separator on Western Farms.
202. Experiment Station Work—XXVI.
203. Canned Fruits, Preserves, and Jellies.
204. The Cultivation of Mushrooms.
205. Pig Management.
206. Milk Fever and Its Treatment.
209. Controlling the Boll Weevil in Cotton Seed and at Ginneries.
210. Experiment Station Work—XXVII.
213. Raspberries.
218. The School Garden.
219. Lessons from the Grain Rust Epidemic of 1904.
220. Tomatoes.
221. Fungous Diseases of the Cranberry.
222. Experiment Station Work—XXVIII.
223. Miscellaneous Cotton Insects in Texas.
224. Canadian Field Peas.
225. Experiment Station Work—XXIX.
227. Experiment Station Work—XXX.
228. Forest Planting and Farm Management.
229. The Production of Good Seed Corn.
231. Spraying for Cucumber and Melon Diseases.
232. Okra: Its Culture and Uses.
233. Experiment Station Work—XXXI.
234. The Guinea Fowl.
235. Preparation of Cement Concrete.
236. Incubation and Incubators.
237. Experiment Station Work—XXXII.
238. Citrus Fruit Growing in the Gulf States.
239. The Corrosion of Fence Wire.
241. Butter Making on the Farm.
242. An Example of Model Farming.
243. Fungicides and Their Use in Preventing Diseases of Fruits.
244. Experiment Station Work—XXXIII.



XUM

245. Renovation of Worn-out Soils
246. Saccharine Sorghums for Forage.
248. The Lawn.
249. Cereal Breakfast Foods.
250. The Prevention of Stinking Smut of Wheat and Loose Smut of Oats.
251. Experiment Station Work—XXXIV.
252. Maple Sugar and Sirup.
253. The Germination of Seed Corn.
254. Cucumbers.
255. The Home Vegetable Garden.
256. Preparation of Vegetables for the Table.
257. Soil Fertility.
258. Texas or Tick Fever and Its Prevention.
259. Experiment Station Work—XXXV.
260. Seed of Red Clover and Its Impurities.
262. Experiment Station Work—XXXVI.
263. Practical Information for Beginners in Irrigation.
264. The Brown-tail Moth and How to Control It.
266. Management of Soils to Conserve Moisture.
267. Experiment Station Work—XXXVII.
268. Industrial Alcohol: Sources and Manufacture.
269. Industrial Alcohol: Uses and Statistics.
270. Modern Conveniences for the Farm Home.
271. Forage Crop Practices in Western Oregon and Western Washington.
272. A Successful Hog and Seed-corn Farm.
273. Experiment Station Work—XXXVIII.
274. Flax Culture.
275. The Gipsy Moth and How to Control It.
276. Experiment Station Work—XXXIX.
277. The Use of Alcohol and Gasoline in Farm Engines.
278. Leguminous Crops for Green Manuring.
279. A Method of Eradicating Johnson Grass.
280. A Profitable Tenant Dairy Farm.
281. Experiment Station Work—XL.
282. Celery.
283. Spraying for Apple Diseases and the Codling Moth in the Ozarks.
284. Insect and Fungous Enemies of the Grape East of the Rocky Mountains.
285. The Advantage of Planting Heavy Cotton Seed.
286. Comparative Value of Whole Cotton Seed and Cotton-seed Meal in Fertilizing Cotton.
287. Poultry Management.
288. Nonsaccharine Sorghums.
289. Beans.
290. The Cotton Bollworm.
291. Evaporation of Apples.
292. Cost of Filling Silos.
293. Use of Fruit as Food.
294. Farm Practice in the Columbia Basin Uplands.
295. Potatoes and Other Root Crops as Food.
296. Experiment Station Work—XLI.
298. Food Value of Corn and Corn Products.
299. Diversified Farming Under the Plantation System.
300. Some Important Grasses and Forage Plants for the Gulf Coast Region.
301. Home-grown Tea.
302. Sea Island Cotton: Its Culture, Improvement, and Diseases.
303. Corn Harvesting Machinery.
304. Growing and Curing Hops.
305. Experiment Station Work—XLII.
306. Dodder in Relation to Farm Seeds.
307. Roselle: Its Culture and Uses.
309. Experiment Station Work—XLIII.
310. A Successful Alabama Diversification Farm.
311. Sand-clay and Burnt-clay Roads.
312. A Successful Southern Hay Farm.
313. Harvesting and Storing Corn.
314. A Method of Breeding Early Cotton to Escape Boll-weevil Damage.
315. Progress in Legume Inoculation.
316. Experiment Station Work—XLIV.
317. Experiment Station Work—XLV.
318. Cowpeas.
319. Demonstration Work in Cooperation with Southern Farmers.
320. Experiment Station Work—XLVI.
321. The Use of the Split-log Drag on Earth Roads.
322. Milo as a Dry-land Grain Crop.
323. Clover Farming on the Sandy Jack-pine Lands of the North.
324. Sweet Potatoes.
325. Small Farms in the Corn Belt.
326. Building Up a Run-down Cotton Plantation.
328. Silver Fox Farming.
329. Experiment Station Work—XLVII.
330. Deer Farming in the United States.
331. Forage Crops for Hogs in Kansas and Oklahoma.
332. Nuts and Their Uses as Food.
333. Cotton Wilt.
334. Experiment Station Work—XLVIII.
335. Harmful and Beneficial Mammals of the Arid Interior.
337. Cropping Systems for New England Dairy Farms.
338. Macadam Roads.
339. Alfalfa.
341. The Basket Willow.
342. Experiment Station Work—XLIX.
343. The Cultivation of Tobacco in Kentucky and Tennessee.
344. The Boll Weevil Problem, with Special Reference to Means of Reducing Damage.
345. Some Common Disinfectants.
346. The Computation of Rations for Farm Animals by the Use of Energy Values.
347. The Repair of Farm Equipment.
348. Bacteria in Milk.
349. The Dairy Industry in the South.
350. The Dehorning of Cattle.
351. The Tuberculin Test of Cattle for Tuberculosis.
352. The Nevada Mouse Plague of 1907–8.
353. Experiment Station Work—L.
354. Onion Culture.
355. A Successful Poultry and Dairy Farm.
356. Peanuts.
357. Methods of Poultry Management at the Maine Agricultural Experiment Station.
358. A Primer of Forestry. Part II: Practical Forestry.
359. Canning Vegetables in the Home.
360. Experiment Station Work—LI.
361. Meadow Fescue: Its Culture and Uses.
362. Conditions Affecting the Value of Market Hay.
363. The Use of Milk as Food.
364. A Profitable Cotton Farm.
365. Farm Management in Northern Potato-growing Sections.
366. Experiment Station Work—LII.
367. Lightning and Lightning Conductors.
368. The Eradication of Bindweed, or Wild Morning-glory
369. How to Destroy Rats.
370. Replanning a Farm for Profit.
371. Drainage of Irrigated Lands.
372. Soy Beans.
373. Irrigation of Alfalfa.
374. Experiment Station Work—LIII.
375. Care of Food in the Home.
376. Game Laws for 1909.
377. Harmfulness of Headache Mixtures.
378. Methods of Exterminating the Texas-fever Tick.
379. Hog Cholera.
380. The Loco-weed Disease.
381. Experiment Station Work—LIV.
382. The Adulteration of Forage-plant Seeds.
383. How to Destroy English Sparrows.
384. Experiment Station Work—LV.
385. Boys' and Girls' Agricultural Clubs.
386. Potato Culture on Irrigated Farms of the West.
387. The Preservative Treatment of Farm Timbers.
388. Experiment Station Work—LVI.
389. Bread and Bread Making.
390. Pheasant Raising in the United States.
391. Economical Use of Meat in the Home.
392. Irrigation of Sugar Beets.
393. Habit-forming Agents.
394. The Use of Windmills in Irrigation in the Semiarid West.
395. Sixty-day and Kherson Oats.
396. The Muskrat.
397. Bees.
398. Farm Practice in the Use of Commercial Fertilizers in the South Atlantic States.
399. Irrigation of Grain.

O